

ROOTS OF DEHN TWISTS ABOUT MULTICURVES

KASHYAP RAJEEVSARATHY AND PRAHLAD VAIDYANATHAN

ABSTRACT. A *multicurve* $\mathcal{C}$ on a closed orientable surface is defined to be a finite collection of disjoint non-isotopic essential simple closed curves. The Dehn twist $t_{\mathcal{C}}$ about $\mathcal{C}$ is the product of the Dehn twists about the individual curves. In this paper, we give necessary and sufficient conditions for the existence of a root of such a Dehn twist, that is, a homeomorphism h such that $h^n = t_{\mathcal{C}}$. We give combinatorial data that corresponds to such roots, and use it to determine upper bounds for n . Finally, we classify all such roots up to conjugacy for surfaces of genus 3 and 4.

1. INTRODUCTION

For $g \geq 0$, let S_g be the closed, orientable surface of genus g , and let $\text{Mod}(S_g)$ denote the mapping class group of S_g . By a *multicurve* $\mathcal{C}$ in S_g , we mean a finite collection of disjoint non-isotopic essential simple closed curves in S_g . Let t_c denote the left-handed Dehn twist about an essential simple closed curve c on S_g . Since the Dehn twists about any two curves in $\mathcal{C}$ commute, we will define the *left-handed Dehn twist about $\mathcal{C}$* to be

$$t_{\mathcal{C}} := \prod_{c \in \mathcal{C}} t_c$$

A *root of $t_{\mathcal{C}}$ of degree n* is an element $h \in \text{Mod}(S_g)$ such that $h^n = t_{\mathcal{C}}$.

When $\mathcal{C}$ comprises a single nonseparating curve, D. Margalit and S. Schleimer [6] showed the existence of roots of $t_{\mathcal{C}}$ of degree $2g - 1$ in $\text{Mod}(S_g)$, for $g \geq 2$. This motivated [7], in which D. McCullough and the first author derived necessary and sufficient conditions for the existence of a root of degree n . As immediate applications of the main theorem in the paper, they showed that roots of even degree cannot exist and that $n \leq 2g - 1$. When $\mathcal{C}$ consists of a single separating curve, the first author derived conditions [9] for the existence of a root of $t_{\mathcal{C}}$. A stable quadratic upper bound on n , and complete classifications of roots for S_2 and S_3 , were derived as corollaries to the main result. In this paper, we shall derive conditions for the existence of a root of $t_{\mathcal{C}}$ when $|\mathcal{C}| \geq 2$, and since there are no such multicurves in S_1 or S_0 , we shall assume henceforth that $g \geq 2$.

In general, a root h of $t_{\mathcal{C}}$ may permute some curves in $\mathcal{C}$, while preserving other curves. So we define an (r, k) -*permuting root* of $t_{\mathcal{C}}$ to be one that

2000 *Mathematics Subject Classification.* Primary 57M99; Secondary 57M60.

Key words and phrases. surface, mapping class, Dehn twist, multicurve, root.

induces a partition of $\mathcal{C}$ into r singletons and k other subsets of size greater than one. The theory for $(r, 0)$ -permuting roots, as we will see, can be obtained by generalizing the theories developed in [7] and [9], which involved the analysis of the fixed point data of finite cyclic actions.

The theory that we intend to develop for (r, k) -permuting roots when $k > 0$ can be motivated by the following example. Consider the multicurve $\mathcal{C}$ in S_5 shown in Figure 1.





FIGURE 1. Nonseparating multicurve of size 5 in S_5

It is apparent that the rotation of S_5 by $2\pi/5$ composed with t_c for some fixed $c \in \mathcal{C}$ is a 5^{th} root of $t_{\mathcal{C}}$ in $\text{Mod}(S_5)$. This is a simple example of a $(0, 1)$ -permuting root, which is obtained by removing invariant disks around pairs of points in two distinct orbits of the $2\pi/5$ rotation of S_0 , and then attaching five 1-handles with full twists. This example indicates that a classification of such roots would require the examination of the orbit information of finite cyclic actions, in addition to their fixed point data. This is a significant departure from existing theories developed in [7] and [9].

Any subset of a multicurve will be called a *submulticurve*. A multicurve $\mathcal{C}$ in S_g is said to be *pseudo-nonseparating* if $\mathcal{C}$ separates S_g , but no proper submulticurve of $\mathcal{C}$ separates S_g . A multicurve that contains no pseudo-nonseparating submulticurves will be called a *nonseparating multicurve*, while a multicurve which is a disjoint union of pseudo-nonseparating multicurves will be called a *separating multicurve*. A multicurve that is neither separating nor nonseparating will be called a *mixed multicurve*. In Figure 2 below, the collection of curves $\mathcal{C} = \{c_1, c_2, c_3, c_4\}$ is a mixed multicurve, while the subcollections $\{c_2, c_3\}$, $\{c_1, c_2, c_3\}$ and $\{c_2, c_4\}$ form pseudo-nonseparating, separating and nonseparating multicurves, respectively.

We start by generalizing the notion of a nested (n, ℓ) -action from [9] to a *permuting (n, r, k) -action*. These are C_n -actions on S_g that have r distinguished fixed points, and k distinguished non-trivial orbits. In Section 3, we introduce the notion of a *permuting data set*, which is a generalization

FIGURE 2. The surface S_5 with a mixed multicurve

of a data set from [9]. We use Thurston's orbifold theory [11, Chapter 13] in Theorem 3.8 to establish a correspondence between permuting (n, r, k) -actions on S_g and permuting data sets of genus g and degree n . In other words, permuting data sets algebraically encode these permuting actions and contain all the relevant orbit and fixed-point information required to classify the roots that will be constructed from these actions.

Let $S_g(\mathcal{C})$ denote the surface obtained from S_g by deleting an annular neighbourhood of $\mathcal{C}$ and capping. In Section 4, we prove that conjugacy classes of roots of Dehn twists about nonseparating multicurves correspond to a special subclass of permuting actions on $S_g(\mathcal{C})$. We use this to obtain the following bounds for the degree of such a root

Corollary. *Let $\mathcal{C}$ be a nonseparating multicurve in S_g of size m , and let h be an (r, k) -permuting root of $t_{\mathcal{C}}$ of degree n .*

(i) *If $r \geq 0$, then*

$$n \leq \begin{cases} 4(g - m) + 2 & : g - m \geq 1 \\ g & : g = m. \end{cases}$$

Furthermore, if $g = m$, then this upper bound is realizable.

(ii) *If $r \geq 1$, then n is odd.*

(iii) *If $r = 1$, then $n \leq 2(g - m) + 1$.*

(iv) *If $r \geq 2$, then $n \leq \frac{g - m + r - 1}{r - 1}$.*

If $\mathcal{C}$ is a separating or a mixed multicurve, $S_g(\mathcal{C})$ will have multiple connected components. In order to classify roots of $t_{\mathcal{C}}$ in this case, we will require multiple actions on the various components of $S_g(\mathcal{C})$ which can be put together and extended to a root of $t_{\mathcal{C}}$ on S_g . In Section 5, we will show that this extension will require compatibility of orbits across components of $S_g(\mathcal{C})$ in addition to compatibility of fixed points as in [9]. As an immediate consequence to this theorem, we obtain quadratic bounds for the degree of the root in terms of the genera of the individual components. Furthermore, we obtain a quadratic stable upper bound for the degree of the root as in [9, Theorem 8.14].

In Sections 6 and 7, we use our theory to obtain a complete classification of roots of $t_{\mathcal{C}}$ on surfaces of genus 3 and 4 respectively. We conclude by proving that a root of $t_{\mathcal{C}}$ cannot lie in the Torelli group of S_g , and also

indicate how our results can be extended to classify roots of finite products of powers of commuting Dehn twists. integrated

2. ROOTS AND THEIR INDUCED PARTITIONS

In this section we shall introduce some preliminary notions, which will be used in later sections.

Notation 2.1. Let $\mathcal{C}$ be a multicurve in S_g , and let N be a closed annular neighbourhood of $\mathcal{C}$.

- (i) We denote the surface $\overline{S_g \setminus N}$ by $\widehat{S_g(\mathcal{C})}$.
- (ii) The closed orientable surface obtained from $\widehat{S_g(\mathcal{C})}$ by capping off its boundary components is denoted by $S_g(\mathcal{C})$.
- (iii) If $\mathcal{C}$ is a nonseparating multicurve, then $S_g(\mathcal{C})$ is a connected surface whose genus we denote by $g_{\mathcal{C}}$.

Notation 2.2. Recall that a multicurve $\mathcal{C}$ in S_g is said to be *pseudo-nonseparating* if $S_g(\mathcal{C})$ is disconnected, but $S_g(\mathcal{C}')$ is not disconnected for any proper submulticurve $\mathcal{C}' \subset \mathcal{C}$.

- (i) If $|\mathcal{C}| = k$, we write $\mathcal{C}^{(k)}$ for such a multicurve. Note that $\mathcal{C}^{(1)}$ is a single separating curve.
- (ii) A disjoint union of m copies of $\mathcal{C}^{(k)}$ is denoted by $\mathcal{C}^{(k)}(m)$.
- (iii) For integers $g \geq 0$ and $m \geq 1$, we define $\mathbb{S}_g(m)$ to be the disjoint union of m copies $\{S_g^1, S_g^2, \dots, S_g^m\}$ of S_g isometrically imbedded in $\mathbb{R}^3$. In particular, $\mathbb{S}_g(1) \cong S_g$, and hence we shall write S_g for $\mathbb{S}_g(1)$.
- (iv) Given two surfaces S_{g_1} and $\mathbb{S}_{g_2}(m)$ and a fixed $k \in \mathbb{N}$, we construct a new surface S_g with $g = (g_1 + mg_2 + (k-1)m)$, containing a multicurve of type $\mathcal{C}^{(k)}(m)$, in the following manner. We remove km disks $\{D_{i,j}^1 : 1 \leq j \leq k, 1 \leq i \leq m\}$ on S_{g_1} and k disks $\{D_{i,j}^2 : 1 \leq j \leq k\}$ on each $S_{g_2}^i$. Now connect $\partial D_{i,j}^1$ to $\partial D_{i,j}^2$ along a 1-handle $A_{i,j}$, and choose the unique curve (upto isotopy) $c_{i,j}$ on $A_{i,j}$. Let $\mathcal{C} = \{c_{i,j}\}$, then note that $\mathcal{C} = \mathcal{C}^{(k)}(m)$, so we write

$$S_{g_1} \#_{\mathcal{C}} \mathbb{S}_{g_2}(m)$$

for the new surface S_g .

- (v) Similarly, given surfaces $\{S_{g_1}, \mathbb{S}_{g_2,1}(m_1), \dots, \mathbb{S}_{g_2,s}(m_s)\}$ and non-negative integers $\{k_1, k_2, \dots, k_s\}$, we construct a new surface S_g with $g = g_1 + \sum_{i=1}^s m_i(g_2 + k_i - 1)$, containing a multicurve of type $\mathcal{C} = \sqcup_{i=1}^s \mathcal{C}^{(k_i)}(m_i)$. Let

$$S_{g'_i} := S_{g_1} \#_{\mathcal{C}^{(k_i)}(m_i)} \mathbb{S}_{g_2,i}(m_i)$$

and $\mathcal{C}_i := \mathcal{C} \setminus \mathcal{C}^{(k_i)}(m_i)$, we now define

$$\overline{\#}_{i=1}^s \left(S_{g_1} \#_{\mathcal{C}^{(k_i)}(m_i)} \mathbb{S}_{g_2,i}(m_i) \right) := \bigcup_{i=1}^s \widehat{S_{g'_i}(\mathcal{C}_i)}.$$

If $s = 2$, we simply write $S_g = \mathbb{S}_{g_2,1}(m_1) \#_{\mathcal{C}^{(k_1)}(m_1)} S_{g_1} \#_{\mathcal{C}^{(k_2)}(m_2)} \mathbb{S}_{g_2,2}(m_2)$.

In Figure 3 below, we give an example of a such a surface S_{22} with a multicurve $\mathcal{C} = \mathcal{C}^{(2)}(2) \sqcup \mathcal{C}^{(1)}(3)$ constructed as in Notation 2.2 from the surfaces $S_5, \mathbb{S}_3(2)$, and $\mathbb{S}_3(3)$.




FIGURE 3. The surface $S_{22} = \mathbb{S}_3(2) \#_{c_1} S_5 \#_{c_2} \mathbb{S}_3(3)$, where $\mathcal{C}_1 = \mathcal{C}^{(2)}(2)$ and $\mathcal{C}_2 = \mathcal{C}^{(1)}(3)$.

Remark 2.3. Let $\mathcal{C} = \{c_1, c_2, \dots, c_m\}$, and suppose that h is a root of $t_{\mathcal{C}}$ of degree n in $\text{Mod}(S_g)$. Then, we claim that, up to isotopy, $h(\mathcal{C}) = \mathcal{C}$.

Suppose first that $h(c_i) \approx c_1$ for all i . Then there exists a neighbourhood N of c_1 such that

$$t_{h(c_i)}|_N = \text{id}_N \text{ and } t_{c_i}|_N = \text{id}_N \quad \forall i \neq 1$$

However, since $h^n = t_{\mathcal{C}}$, it follows that

$$t_{\mathcal{C}} = ht_{\mathcal{C}}h^{-1} = ht_{c_1}h^{-1}ht_{c_2}h^{-1} \dots ht_{c_m}h^{-1} = t_{h(c_1)}t_{h(c_2)} \dots t_{h(c_m)}$$

Hence

$$t_{c_1}|_N = t_{\mathcal{C}}|_N = t_{h(c_1)}t_{h(c_2)} \dots t_{h(c_m)}|_N = \text{id}_N$$

which is a contradiction.

So we assume without loss of generality that $h(c_1) \sim c_1$. Then we choose a neighbourhood N' of c_1 disjoint from c_i for $i \neq 1$, and choose an isotopy φ_t such that

$$\varphi_t|_{S_g \setminus N'} = \text{id}_{S_g \setminus N'}$$

and $\varphi_0 = h$ and $\varphi_1(c_1) = c_1$. Replacing h by φ_1 , we may assume that $h(c_1) = c_1$. Now note that $h(c_i) \approx c_1$ for all $i \neq 1$, which allows us to proceed by induction on $|\mathcal{C}|$ to conclude that, up to isotopy, $h(\mathcal{C}) = \mathcal{C}$.

Definition 2.4. Let $\mathcal{C}$ be a multicurve of size m in S_g . Then for integers $r, k \geq 0$, an (r, k) -partition of $\mathcal{C}$ is a partition $\mathbb{P}_{r,k}(\mathcal{C}) = \{\mathcal{C}'_1, \dots, \mathcal{C}'_r, \mathcal{C}_1, \dots, \mathcal{C}_k\}$ of the set $\mathcal{C}$ into subsets such that for all i ,

- (i) $|\mathcal{C}'_i| = 1$, $|\mathcal{C}_i| > 1$, and
- (ii) $\mathcal{C}_i$ comprises only separating or only nonseparating curves.

Definition 2.5. Let $\mathcal{C}$ be a multicurve in S_g . Then for integers $r, k \geq 0$, a root h of $t_{\mathcal{C}}$ of is said to be (r, k) -permuting if it induces an (r, k) -partition of $\mathcal{C}$.

Notation 2.6. Let $\mathcal{C}$ be a multicurve of size m in S_g and consider an (r, k) -partition $\mathbb{P}_{r,k}(\mathcal{C}) = \{\mathcal{C}'_1, \dots, \mathcal{C}'_r, \mathcal{C}_1, \dots, \mathcal{C}_k\}$ as in Definition 2.4.

- (i) We shall denote the multiset $\{\{|\mathcal{C}_1|, |\mathcal{C}_2|, \dots, |\mathcal{C}_k|\}\}$ by $S(\mathbb{P}_{r,k})$. (From here on, we shall denote a multiset using $\{\{ \} \}$).
- (ii) If $\mathbb{P}_{r,k}(\mathcal{C})$ is induced by an (r, k) -permuting root h of $t_{\mathcal{C}}$, then we shall denote it by $\mathcal{C}_{r,k}(h)$.

3. PERMUTING ACTIONS AND PERMUTING DATA SETS

In this section we shall introduce permuting (n, r, k) -actions, which are generalizations of the nested (n, ℓ) -actions from [9]. We shall also introduce the notion of a permuting (n, r, k) -data set, which is an abstract tuple involving non-negative integers that would algebraically encode a permuting (n, r, k) -action.

Definition 3.1. For integers $n \geq 1$ and $r, k \geq 0$, an orientation-preserving C_n -action t on S_g is called a *permuting (n, r, k) -action* if

- (i) there is a set $\mathbb{P}(t) \subset S_g$ of r distinguished fixed points of t , and
- (ii) there is a set $\mathbb{O}(t) \subset S_g$ of k distinguished non-trivial orbits of t .

Notation 3.2. Let t be a permuting (n, r, k) -action on S_g .

- (i) Fix a point $P \in S_g$, and consider $t_* : T_P(S_g) \rightarrow T_{t(P)}(S_g)$. By the Nielsen realisation theorem [5], we may change t by isotopy in $\text{Mod}(S_g)$ so that t_* is an isometry. Hence, t_* induces a local rotation by an angle, which we shall denote by $\theta_P(t)$. Note that if $P \in \mathbb{P}(t)$, then $\theta_P(t) = 2\pi a/n$, where $\gcd(a, n) = 1$.
- (ii) Fix an orbit $\mathbb{O} = \{P_1, \dots, P_s\} \in \mathbb{O}(t)$. If $s < n$, then $s \mid n$, and there exists a cone point in the quotient orbifold of degree n/s . Each P_i has stabilizer generated by t^s and the rotation induced by t^s around each P_i must be the same, since its action at one point is conjugate by a power of t to its action at each other point in the orbit. So the rotation angle is of the form $2\pi c^{-1}/(n/s) \pmod{2\pi}$, where $(c, n/s) = 1$ and c^{-1} denotes the inverse of $c \pmod{n/s}$. We now associate to this orbit a pair $p(\mathbb{O})$ as follows:

$$p(\mathbb{O}) := \begin{cases} (c, n/s) & \text{if } s < n \\ (0, 1) & \text{if } s = n. \end{cases}$$

- (iii) For any orbit $\mathbb{O} \in \mathbb{O}(t)$, if $p(\mathbb{O}) = (a, b)$, then we define

$$\theta_{\mathbb{O}}(t) := \begin{cases} 2\pi a^{-1}/b & \text{if } a \neq 0 \\ 0 & \text{if } a = 0. \end{cases}$$

Definition 3.3. Consider a permuting (n, r, k) -action t on S_g with $\mathbb{P}(t) = \{P_1, \dots, P_r\}$ and $\mathbb{O}(t) = \{\mathbb{O}_1, \mathbb{O}_2, \dots, \mathbb{O}_k\}$.

- (i) We write

$$S(t) = \{\{|\mathbb{O}_1|, |\mathbb{O}_2|, \dots, |\mathbb{O}_k|\}\}.$$

(ii) For each $p \in \{p(\mathbb{O}_i) : 1 \leq i \leq k\}$, define

$$m_p = |\{j : p(\mathbb{O}_j) = x\}|.$$

We define the *orbit distribution* of t to be the set

$$\mathbb{O}_t = \{(p, m_p) : p \in \{p(\mathbb{O}_i) : 1 \leq i \leq k\}\}.$$

Definition 3.4. Let t_1 and t_2 be two permuting (n, r, k) -actions on S_g with $\mathbb{P}(t_s) = \{P_{s,1}, P_{s,2}, \dots, P_{s,r}\}$ and $\mathbb{O}(t_s) = \{\mathbb{O}_{s,1}, \mathbb{O}_{s,2}, \dots, \mathbb{O}_{s,k}\}$ for $s = 1, 2$. We say t_1 is *equivalent* to t_2 if $\mathbb{O}_{t_1} = \mathbb{O}_{t_2}$ and there is an orientation-preserving homeomorphism $\phi \in \text{Mod}(S_g)$ such that

- (i) $\phi(P_{1,i}) = P_{2,i}$ for $1 \leq i \leq r$,
- (ii) for each $1 \leq j \leq k$, if $\mathbb{O}_{s,j} = \{Q_{j,1}^s, Q_{j,2}^s, \dots, Q_{j,m_{s,j}}^s\}$, then $m_{1,j} = m_{2,j}$ and $\phi(Q_{j,i}^1) = Q_{j,i}^2$ for all $1 \leq i \leq m_{1,j}$, and
- (iii) $\phi t_1 \phi^{-1}$ is isotopic to t_2 relative to $\mathbb{P}(t_2) \sqcup (\cup_{j=1}^k \mathbb{O}_{2,j})$.

The equivalence class of a permuting (n, r, k) -action is denoted by $\llbracket t \rrbracket$.

We now introduce the notion of an (n, r) -*data set*, which encodes the signature of the quotient orbifold of a permuting (n, r, k) -action and the turning angles around its distinguished fixed points. Furthermore, the (n, r) -data set will be combined with the orbit distribution of the action to form a pair, which we will call a *permuting (n, r, k) -data set*.

Definition 3.5. Given $n \geq 1$ and $r \geq 0$, an (n, r) -*data set* is a tuple

$$\mathcal{D} = (n, g_0, (a_1, a_2, \dots, a_r); (c_1, n_1), (c_2, n_2), \dots, (c_s, n_s))$$

where $n \geq 1$ and $g_0 \geq 0$ are integers, each a_i is a residue class modulo n , and each c_i is a residue class modulo n_i such that:

- (i) each $n_i \mid n$,
- (ii) $\gcd(a_i, n) = \gcd(c_i, n_i) = 1$, and
- (iii) $\sum_{i=1}^r a_i + \sum_{j=1}^s \frac{n}{n_j} c_j \equiv 0 \pmod{n}$.

The number g determined by the equation

$$(1) \quad \frac{2-2g}{n} = 2 - 2g_0 + r \left(\frac{1}{n} - 1 \right) + \sum_{j=1}^s \left(\frac{1}{n_j} - 1 \right)$$

is called the *genus* of the data set.

Definition 3.6. Fix an (n, r) -data set $\mathcal{D}$ of genus g as above.

- (i) For each $(a, b) \in \{(0, 1), (c_1, n_1), \dots, (c_s, n_s)\}$, we write

$$\theta((a, b)) := \begin{cases} 0 & \text{if } a = 0, \\ 2\pi a^{-1}/b & \text{if } a \neq 0. \end{cases}$$

- (ii) For each $p \in \{(0, 1), (c_1, n_1), \dots, (c_s, n_s)\}$, choose a non-negative integer m_p . Then the set $\mathbb{O}_{\mathcal{D}} = \{(p, m_p) : m_p > 0\}$ is called an *orbit distribution* of $\mathcal{D}$.

- (iii) Given an orbit distribution $\mathbb{O}_{\mathcal{D}}$ associated with an (n, r) -data set $\mathcal{D}$, the pair $(\mathcal{D}, \mathbb{O}_{\mathcal{D}})$ is called a *permuting (n, r, k) -data set* of genus g , where $k = \sum_p m_p$.

Definition 3.7. Let

$$\mathcal{D} = (n, g_0, (a_1, a_2, \dots, a_r); (c_1, n_1), (c_2, n_2), \dots, (c_s, n_s))$$

$$\text{and } \mathcal{D}' = (n, g'_0, (a'_1, a'_2, \dots, a'_r); (c'_1, n'_1), (c'_2, n'_2), \dots, (c'_s, n'_s))$$

be two (n, r) -data sets as in Definition 3.5.

- (i) $\mathcal{D}$ and $\mathcal{D}'$ are said to be *equivalent* if

$$\{\{a_1, a_2, \dots, a_r\}\} = \{\{a'_1, a'_2, \dots, a'_r\}\}, \text{ and}$$

$$\{\{(c_1, n_1), \dots, (c_s, n_s)\}\} = \{\{(c'_1, n'_1), \dots, (c'_s, n'_s)\}\}.$$

- (ii) Two permuting (n, r, k) -data sets $(\mathcal{D}, \mathbb{O}_{\mathcal{D}})$ and $(\mathcal{D}', \mathbb{O}_{\mathcal{D}'})$ are said to be *equivalent* if $\mathcal{D}$ and $\mathcal{D}'$ are equivalent as above, and $\mathbb{O}_{\mathcal{D}} = \mathbb{O}_{\mathcal{D}'}$.

Note that equivalent data sets have the same genus.

Theorem 3.8. *Given $n \geq 1$ and $g \geq 0$, equivalence classes of permuting (n, r, k) -data sets of genus g correspond to equivalence classes of permuting (n, r, k) -actions on S_g .*

Proof. Let t be a permuting (n, r, k) -action on S_g with quotient orbifold $\mathcal{O}$ whose underlying surface has genus g_0 . If t is a free action, then $\mathcal{O} = S_{g_0}$, and we simply write $\mathcal{D} = (n, g_0;)$ and $\mathbb{O}_{\mathcal{D}} = \{(0, 1), k\}$. If t is not free, let p_j be the image in $\mathcal{O}$ of the P_j , for $1 \leq j \leq r$, and let $q_1, q_2, \dots, q_s$ be the other possible cone points of $\mathcal{O}$ as in Figure 3.




FIGURE 4. The quotient orbifold $\mathcal{O}$.

Let α_i be the generator of the orbifold fundamental group $\pi_1^{orb}(\mathcal{O})$ that goes around the point p_i , $1 \leq i \leq r$ and let γ_j be the generators going around q_j , $1 \leq j \leq s$. Let x_p and y_p , $1 \leq p \leq g_0$ be the standard generators of the “surface part” of $\mathcal{O}$, chosen to give the following presentation of $\pi_1^{orb}(\mathcal{O})$:

$$\pi_1^{orb}(\mathcal{O}) = \langle \alpha_1, \alpha_2, \dots, \alpha_r, \gamma_1, \gamma_2, \dots, \gamma_s, x_1, y_1, x_2, y_2, \dots, x_{g_0}, y_{g_0} \mid$$

$$\alpha_1^n = \alpha_2^n = \dots = \gamma_1^{n_1} = \gamma_2^{n_2} = \dots = \gamma_s^{n_s} = 1, \alpha_1 \dots \alpha_r \gamma_1 \dots \gamma_s = \prod_{p=1}^{g_0} [x_p, y_p] \rangle$$

From orbifold covering space theory [11], we have the following exact sequence

$$1 \rightarrow \pi_1(S_g) \rightarrow \pi_1^{orb}(\mathcal{O}) \xrightarrow{\rho} C_n \rightarrow 1,$$

where $C_n = \langle t \rangle$. The homomorphism ρ is obtained by lifting path representatives of elements of $\pi_1^{orb}(\mathcal{O})$. Since these do not pass through the cone points, the lifts are uniquely determined.

For $1 \leq i \leq s$, the preimage of q_i consists of n/n_i points cyclically permuted by t . As in Notation 3.2, the rotation angle at each point is of the form $2\pi c_i^{-1}/n_i$ where c_i is a residue class modulo n_i and $\gcd(c_i, n_i) = 1$. Lifting the γ_i , we have that $\rho(\gamma_i) = h^{(n/n_i)c_i}$. Similarly, lifting the α_i gives $\rho(\alpha_i) = t^{a_i}$ where $\gcd(a_i, n) = 1$. Finally, we have

$$\rho\left(\prod_{p=1}^{g_0} [x_p, y_p]\right) = 1,$$

since C_n is abelian, so

$$1 = \rho(\alpha_1 \dots \alpha_r \gamma_1 \dots \gamma_s) = t^{a_1 + \dots + a_r + (n/n_1)c_1 + \dots + (n/n_s)c_s}$$

giving

$$\sum_{i=1}^r a_i + \sum_{j=1}^s \frac{n}{n_j} c_j \equiv 0 \pmod{n}.$$

The fact that the data set $\mathcal{D}$ has genus g follows easily from the multiplicativity of the orbifold Euler characteristic for the orbifold covering $S_g \rightarrow \mathcal{O}$:

$$\frac{2-2g}{n} = 2 - 2g_0 + r \left(\frac{1}{n} - 1 \right) + \sum_{j=1}^s \left(\frac{1}{n_j} - 1 \right).$$

Thus, h gives a (n, r) -data set

$$\mathcal{D} = (n, g_0, (a_1, a_2, \dots, a_r); (c_1, n_1), (c_2, n_2), \dots, (c_s, n_s))$$

of genus g , and hence $(\mathcal{D}, \mathbb{O}_t)$ forms a permuting (n, r, k) -data set.

Consider another permuting (n, r, k) -action t' in the equivalence class of t with a distinguished fixed point set $\mathbb{P}(t') = \{P'_1, P'_2, \dots, P'_r\}$. Then by definition there exists an orientation-preserving homeomorphism $\phi \in \text{Mod}(S_g)$ such that $\phi(P_j) = P'_j$ for all j and $\phi t \phi^{-1}$ is isotopic to t' relative to $\mathbb{P}(t')$. Therefore, $\theta_{P_j}(t) = \theta_{P'_j}(t')$, for $1 \leq j \leq r$, and since $\mathbb{O}_t = \mathbb{O}_{t'}$, the two actions will produce the same permuting (n, r, k) -data sets.

Conversely, given a permuting (n, r, k) -data set $(\mathcal{D}, \mathbb{O}_{\mathcal{D}})$, we construct the orbifold $\mathcal{O}$ and a representation $\rho : \pi_1^{orb}(\mathcal{O}) \rightarrow C_n$. Any finite subgroup of $\pi_1^{orb}(\mathcal{O})$ is conjugate to one of the cyclic subgroups generated by α_j or a γ_i , so condition (iv) in the definition of the data set ensures that the kernel of ρ is torsion-free. Therefore, the orbifold covering $S \rightarrow \mathcal{O}$ corresponding to the kernel is a manifold, and calculation of the Euler characteristic shows that $S = S_g$. Thus we obtain a C_n -action t on S_g with r distinguished fixed points $\mathbb{P}(t)$. We now construct $\mathbb{O}_t$ from $\mathbb{O}_{\mathcal{D}}$ in the following manner. For

each pair $(p, m_p) \in \mathbb{O}_{\mathcal{D}}$, write $p = (a, b)$. If $a = 0$, then choose m_p orbits of size n (if t is a free action, this choice is trivial, but otherwise, such an orbit would always exist in a small neighbourhood around any fixed point of t). If $a \neq 0$, then there exists a cone point in $\mathcal{O}$ of degree b , so there exists an orbit of t of size n/b in S_g . Once again, by considering a small neighbourhood of this orbit, we may choose m_p distinct orbits $\{\mathbb{O}_p^1, \mathbb{O}_p^2, \dots, \mathbb{O}_p^{m_p}\}$ and set

$$\mathbb{O}(t) := \bigsqcup_{(p, m_p) \in \mathbb{O}_{\mathcal{D}}} \{\mathbb{O}_p^1, \mathbb{O}_p^2, \dots, \mathbb{O}_p^{m_p}\},$$

which in turn gives $\mathbb{O}_t = \mathbb{O}_{\mathcal{D}}$.

It remains to show that the resulting action on S_g is determined upto our equivalence in $\text{Mod}(S_g)$. Suppose that two permuting (n, r, k) -actions t and t' have the same permuting (n, r, k) -data set $(\mathcal{D}, \mathbb{O}_{\mathcal{D}})$. $\mathcal{D}$ encodes the fixed point data of the periodic transformation t , so by a result of J. Nielsen [8] (or by a subsequent result of A. Edmonds [2, Theorem 1.3]), t and t' have to be conjugate by an orientation-preserving homeomorphism ϕ . Let $\mathcal{O}'$ be the quotient orbifold of the action t' , and $\rho' : \pi_1^{orb}(\mathcal{O}') \rightarrow C_n$ be the induced representations. Then ϕ induces a map $\varphi_{\#} : \pi_1^{orb}(\mathcal{O}) \rightarrow \pi_1^{orb}(\mathcal{O}')$ such that $\rho' \circ \varphi_{\#} = \rho$ as in [7, Theorem 2.1]. If γ is a loop around a cone point in $\mathcal{O}$, then $\rho(\gamma)$ is a loop around a cone point in $\mathcal{O}'$, and these cone points are associated to the same pair in $\mathcal{D}$ since $\rho'(\varphi_{\#}(\gamma)) = \rho(\gamma)$. Hence, ϕ maps $\mathbb{P}(t)$ to $\mathbb{P}(t')$ and $\mathbb{O}(t)$ to $\mathbb{O}(t')$ as in Definition 3.4. Furthermore, $\mathbb{O}_t = \mathbb{O}_{\mathcal{D}} = \mathbb{O}_{t'}$ by construction, and hence the permuting data set determines t upto equivalence. $\square$

4. NONSEPARATING MULTICURVES

Recall that a multicurve $\mathcal{C}$ is said to be nonseparating if it does not contain any pseudo-nonseparating submulticurves. In this section, we establish that a root of $t_{\mathcal{C}}$ corresponds to a special kind of permuting action on the connected surface $S_g(\mathcal{C})$.

Definition 4.1. Let t_i be a permuting (n_i, r_i, k_i) -action on S_{g_i} for $i = 1, 2$. Two orbits $\mathbb{O}_i \in \mathbb{O}(t_i)$ are said to be *equivalent* (in symbols, $\mathbb{O}_1 \sim \mathbb{O}_2$) if

- (i) $|\mathbb{O}_1| = |\mathbb{O}_2|$, and
- (ii) if $|\mathbb{O}_1| < n := \text{lcm}(n_1, n_2)$, then we further require that

$$\theta_{\mathbb{O}_1}(t_1) + \theta_{\mathbb{O}_2}(t_2) \equiv 2\pi/n \pmod{2\pi}.$$

In this section, we will only need the case when $t_1 = t_2$, but we will need the general case in Section 5.

Definition 4.2. Let $\mathcal{C}$ be a nonseparating multicurve in S_g . A permuting $(n, 2r, 2k)$ -action t on $S_g(\mathcal{C})$ is said to be *nonseparating with respect to $\mathcal{C}$* if there exists a (r, k) -partition $\mathbb{P}_{r,k}(\mathcal{C})$ of $\mathcal{C}$ such that

- (i) there exists r mutually disjoint pairs $\{P_i, P'_i\}$ of distinguished fixed points in $\mathbb{P}(t)$ such that $\theta_{P_i}(t) + \theta_{P'_i}(t) = 2\pi/n$ modulo 2π , for $1 \leq i \leq r$, and

- (ii) there exists k mutually disjoint pairs $\{\mathbb{O}_i, \mathbb{O}'_i\}$ of distinguished nontrivial orbits in $\mathbb{O}_t$ such that $\mathbb{O}_i \sim \mathbb{O}'_i$, for $1 \leq i \leq k$, and
- (iii) $S(\mathbb{P}_{r,k}) = S(t)$.

Theorem 4.3. *Let $\mathcal{C}$ be a nonseparating multicurve in S_g . Then for $n \geq 1$, equivalence classes of permuting $(n, 2r, 2k)$ -actions on $S_g(\mathcal{C})$ that are nonseparating with respect to $\mathcal{C}$ correspond to the conjugacy classes in $\text{Mod}(S_g)$ of (r, k) -permuting roots of $t_{\mathcal{C}}$ of degree n .*

Proof. First, we shall prove that a conjugacy class of an (r, k) -permuting root h of $t_{\mathcal{C}}$ of degree n yields an equivalence class of a permuting $(n, 2r, 2k)$ -action that is nonseparating with respect to $\mathcal{C}$. We assume without loss of generality that $r, k > 0$, with the implicit understanding that, when either of them is zero, the corresponding arguments may be disregarded.

Let $\mathcal{C}_{r,k}(h) = \{\mathcal{C}'_1, \mathcal{C}'_2, \dots, \mathcal{C}'_r, \mathcal{C}_1, \mathcal{C}_2, \dots, \mathcal{C}_k\}$ be the (r, k) -partition of $\mathcal{C}$ associated with h . Choose a closed tubular neighborhood N of $\mathcal{C}$, and consider $S_g(\mathcal{C})$ as in Definition 2.1. By isotopy, we may assume that $t_{\mathcal{C}}(\mathcal{C}) = \mathcal{C}$, $t_{\mathcal{C}}(N) = N$, and $t_{\mathcal{C}}|_{\widehat{S_g(\mathcal{C})}} = \text{id}_{\widehat{S_g(\mathcal{C})}}$. Suppose that h is a root of $t_{\mathcal{C}}$ of degree n , then by Remark 2.3, we may assume that h preserves $\mathcal{C}$ and takes N to N .

By the Nielsen-Kerckhoff theorem [5], $\hat{t} := h|_{\widehat{S_g(\mathcal{C})}}$ is isotopic to a homeomorphism whose n^{th} power is $\text{id}_{\widehat{S_g(\mathcal{C})}}$. So we may change h by isotopy so that $\hat{t}^n = \text{id}_{\widehat{S_g(\mathcal{C})}}$. We fill in the $2m$ boundary circles of $\widehat{S_g(\mathcal{C})}$ with disks and extend $\hat{t}$ to a homeomorphism t on $S_g(\mathcal{C})$ by coning. Thus t defines a effective C_n -action on $S_g(\mathcal{C})$, where $C_n = \langle t \mid t^n = 1 \rangle$.

The C_n -action t fixes the centers P_i and P'_i of the $2r$ disks D_i and D'_i , $1 \leq i \leq r$, of $S_g \setminus \widehat{S_g(\mathcal{C})}$ whose boundaries are the components of ∂N which are fixed by t . The orientation of S_g determines one for $S_g(\mathcal{C})$, so we may speak of directed angles of rotation about the centres of these disks. Since $h^n = t_{\mathcal{C}}$, it follows from [7, Theorem 2.1] that

$$\theta_{P_i}(t) + \theta_{P'_i}(t) = 2\pi/n \pmod{2\pi},$$

as illustrated in Figure 5 below.




FIGURE 5. Angle compatibility at each pair $\{P_i, P'_i\} \subset \mathbb{P}(t)$.

The remaining disks occuring in $\overline{S_g \setminus \widehat{S_g(\mathcal{C})}}$ form k pairs of orbits of sizes $m_1, m_2, \dots, m_k$ where $S(\mathcal{C}_{r,k}(h)) = \{\{m_1, m_2, \dots, m_k\}\}$. For $1 \leq j \leq k$, we denote the centres of these pairs of disks by $Q_{i,j}$ and $Q'_{i,j}$, and the orbits of these centres by $\mathbb{O}_j$ and $\mathbb{O}'_j$. Thus t is a permuting $(n, 2r, 2k)$ -action with $\mathbb{P}(t) = \{P_1, P'_1, \dots, P_r, P'_r\}$ and $\mathbb{O}(t) = \{\mathbb{O}_1, \mathbb{O}'_1, \dots, \mathbb{O}_k, \mathbb{O}'_k\}$.

It remains to show that $\mathbb{O}_i \sim \mathbb{O}'_i$ for each i . By construction, $|\mathbb{O}_i| = |\mathbb{O}'_i| = m_i$, and if $m_i = n$, then $\mathbb{O}_i \sim \mathbb{O}'_i$ holds trivially. If not, then we need to show that the angle compatibility condition from Definition ?? holds. Write $\mathbb{O}_i = \{Q_{i,1}, Q_{i,2}, \dots, Q_{i,m_i}\}$, $\mathbb{O}'_i = \{Q'_{i,1}, Q'_{i,2}, \dots, Q'_{i,m_i}\}$ and note that h^{m_i} is an $(n/m_i)^{\text{th}}$ root of $t_{\mathcal{C}}$ such that $h^{m_i}(c_{i,1}) = c_{i,1}$. Hence,

$$\theta_{Q_{i,1}}(t^{m_i}) + \theta_{Q'_{i,1}}(t^{m_i}) = 2\pi/(n/m_i) \pmod{2\pi},$$

which implies that

$$m_i \theta_{\mathbb{O}_i}(t) + m_i \theta_{\mathbb{O}'_i}(t) \equiv 2\pi/(n/m_i) \pmod{2\pi}.$$

Since t^n is a restriction of a single Dehn twist, it follows that

$$\theta_{\mathbb{O}_i}(t) + \theta_{\mathbb{O}'_i}(t) \equiv 2\pi/n \pmod{2\pi}.$$

Hence $\mathbb{O}_i \sim \mathbb{O}'_i$, and we obtain a permuting $(n, 2r, 2k)$ -action t on $S_g(\mathcal{C})$ that is nonseparating with respect to $\mathcal{C}$.

Now suppose $h_1, h_2 \in \text{Mod}(S_g)$ are two roots of $t_{\mathcal{C}}$ that are conjugate in $\text{Mod}(S_g)$ via $\Phi \in \text{Mod}(S_g)$, and let t_s denote the finite order homeomorphisms on $S_g(\mathcal{C})$ induced by h_s , for $s = 1, 2$. Then $t_{\mathcal{C}} = \phi t_{\mathcal{C}} \Phi^{-1} = t_{\Phi(\mathcal{C})}$, so we may assume upto isotopy that $\Phi(\mathcal{C}) = \mathcal{C}$ (as in Remark 2.3) and that $\Phi(N) = N$. We extend $\Phi|_{\widehat{S_g(\mathcal{C})}}$ to an element $\phi \in \text{Mod}(S_g(\mathcal{C}))$ by coning. Now, ϕ maps $\mathbb{P}(t_1)$ to $\mathbb{P}(t_2)$, and $\mathbb{O}(t_1)$ to $\mathbb{O}(t_2)$ bijectively as in Definition 3.4. Since the h_s and Φ all preserve N , $\phi t_1 \phi^{-1}$ is isotopic to t_2 preserving $\mathbb{P}(t_2)$ and $\mathbb{O}(t_2)$. Furthermore, for each $\mathbb{O} \in \mathbb{O}(t_1)$, $p(\phi(\mathbb{O})) = p(\mathbb{O})$. Hence, $\mathbb{O}_{t_1} = \mathbb{O}_{t_2}$ and so t_1 and t_2 will be equivalent permuting $(n, 2r, 2k)$ -actions.

Conversely, given a permuting $(n, 2r, 2k)$ -action t on $S_g(\mathcal{C})$ that is nonseparating with respect to $\mathcal{C}$, we can reverse the argument to produce the (r, k) -permuting root h . Let $\mathbb{P}(t) = \{P_1, P'_1, \dots, P_r, P'_r\}$ and $\mathbb{O}(t) = \{\mathbb{O}_1, \mathbb{O}'_1, \dots, \mathbb{O}_k, \mathbb{O}'_k\}$. For $1 \leq i \leq r$, we remove disks D_i and D'_i invariant under the action of t around the P_i and P'_i and attaching r annuli to obtain the surface $S_g(\mathcal{C} \setminus \mathcal{C}')$. The condition on the angles $\{\theta_{P_i}(t), \theta_{P'_i}(t)\}$ ensures that the rotation angles work correctly to allow an extension of t to obtain an h_0 with $h_0^n = t_{\mathcal{C}'}$ in $\text{Mod}(S_g(\mathcal{C} \setminus \mathcal{C}'))$, where $\mathcal{C}' = \cup_{i=1}^r \mathcal{C}'_i$.

Now write $\mathbb{O}_1 = \{Q_{1,1}, Q_{1,2}, \dots, Q_{1,m_1}\}$ and consider disks $D_{1,i}$ around $Q_{1,i}$ such that $t(D_{1,i}) = t(D_{1,i+1})$. Similarly, write $\mathbb{O}'_1 = \{Q'_{1,1}, Q'_{1,2}, \dots, Q'_{1,m_1}\}$ and consider disks $D'_{1,i}$ as earlier. Then we attach m_1 annuli connecting $\partial D_{1,i}$ to $\partial D'_{1,i}$. Each such annulus contains a nonseparating curve $c_{1,i}$, which is unique upto isotopy. Repeating this process for $1 \leq i \leq m_1$, we obtain the surface $S_g(\mathcal{C} \setminus (\mathcal{C}' \cup \mathcal{C}_1))$. Since $t(D_{1,i}) = D_{1,i+1}$, we may extend

the homeomorphism h_0 to a homeomorphism $\widetilde{h}_0 \in \text{Mod}(S_g(\mathcal{C}' \cup \mathcal{C}_1))$, which cyclically permutes the $c_{1,i}$. If $|\mathbb{O}_1| = |\mathbb{O}'_1| = n$, then define $h_1 := \widetilde{h}_0 t_{c_{1,1}}$. Otherwise, since $\mathbb{O}_1 \sim \mathbb{O}'_1$, the difference in the turning angles around $Q_{1,i}$ and $Q'_{1,i}$ is $2\pi/n$. Let $\widetilde{h}_1$ be the $(1/n)^{th}$ -twist around $c_{1,1}$. Now $h_1 := \widetilde{h}_0 h_1$ is an $(r, 1)$ -permuting root of $t_{\mathcal{C}' \cup \mathcal{C}_1}$ of degree n in $\text{Mod}(S_g(\mathcal{C} \setminus (\mathcal{C}' \cup \mathcal{C}_1)))$. We now repeat this process inductively to obtain an (r, k) -permuting root $h := h_k \in \text{Mod}(S_g)$ of $t_{\mathcal{C}}$ of degree n .

It remains to show that the resulting root h of $t_{\mathcal{C}}$ is determined up to conjugacy. Suppose t_1 and t_2 are two equivalent $(n, 2r, 2k)$ -actions on $S_g(\mathcal{C})$ that are nonseparating with respect to $\mathcal{C}$ with $\mathbb{P}(t_s) = \{P_{s,1}, P_{s,2}, \dots, P_{s,r}\}$ and $\mathbb{O}(t_s) = \{\mathbb{O}_{s,1}, \mathbb{O}_{s,2}, \dots, \mathbb{O}_{s,k}\}$, for $s = 1, 2$. Let $\phi \in \text{Mod}(S_g(\mathcal{C}))$ be an orientation-preserving homeomorphism satisfying the conditions in Definition 3.4. Then repeating the argument from [7, Theorem 2.1], ϕ extends to a homeomorphism $\Phi_0 \in \text{Mod}(S_g(\mathcal{C} \setminus \mathcal{C}'))$ such that $\Phi_0 h_{1,0} \Phi_0^{-1} = h_{2,0}$, where $h_{s,0}$ is the root of $t_{\mathcal{C}'}$ obtained from t_s , for $s = 1, 2$, as above. Furthermore, since ϕ maps $\mathbb{O}_{1,i}$ to $\mathbb{O}_{2,i}$ as in Definition 3.4, we may once again extend Φ_0 to a homeomorphism $\Phi \in \text{Mod}(S_g)$ satisfying $\Phi h_1 \Phi^{-1} = h_2$, where h_s is the root of $t_{\mathcal{C}}$ obtained from t_s , for $s = 1, 2$. $\square$

Definition 4.4. A permuting $(n, 2r, 2k)$ -data set $(\mathcal{D}, \mathbb{O}_{\mathcal{D}})$ is called *nonseparating* if

- (i) $\mathcal{D} = (n, g_0, (a_1, a'_1, \dots, a_r, a'_r); (c_1, n_1), \dots, (c_s, n_s))$, where $a_i + a'_i \equiv a_i a'_i \pmod{n}$, for $1 \leq i \leq r$
- (ii) For each $(p, m_p) \in \mathbb{O}_{\mathcal{D}}$, there exists $(p', m'_p) \in \mathbb{O}_{\mathcal{D}}$ such that $m_p = m'_p$ and

$$\theta(p) + \theta(p') = \begin{cases} 0 & : p = p' = (0, 1) \\ 2\pi/n \pmod{2\pi} & : \text{otherwise.} \end{cases}$$

We now deduce the following theorem from Theorems 3.8 and 4.3.

Theorem 4.5. Let $\mathcal{C}$ be a nonseparating multicurve in S_g . For $n \geq 1$, equivalence classes of nonseparating permuting $(n, 2r, 2k)$ -data sets of genus $g_{\mathcal{C}}$ correspond to equivalence classes of permuting $(n, 2r, 2k)$ -actions on $S_g(\mathcal{C})$ that are nonseparating with respect to $\mathcal{C}$.

Corollary 4.6. Let $\mathcal{C}$ be a nonseparating multicurve in S_g . For $n \geq 1$, equivalence classes of nonseparating permuting $(n, 2r, 2k)$ -data sets of genus $g_{\mathcal{C}}$ correspond to conjugacy classes in $\text{Mod}(S_g)$ of (r, k) -permuting roots of $t_{\mathcal{C}}$ of degree n .

If $\mathcal{C}$ is a nonseparating multicurve of size m , then $g_{\mathcal{C}} = g - m$. Hence we obtain the following restrictions on the degree of a root of $t_{\mathcal{C}}$.

Corollary 4.7. Let $\mathcal{C}$ be a nonseparating multicurve in S_g of size m , and let h be an (r, k) -permuting root of $t_{\mathcal{C}}$ of degree n .

(i) If $r \geq 0$, then

$$n \leq \begin{cases} 4(g-m) + 2, & \text{if } g-m \geq 1 \\ g, & \text{if } g = m. \end{cases}$$

Furthermore, if $g = m$, then this upper bound is realizable.

(ii) If $r \geq 1$, then n is odd.

(iii) If $r = 1$, then $n \leq 2(g-m) + 1$.

(iv) If $r \geq 2$, then $n \leq \frac{g-m+r-1}{r-1}$.

Proof. If $r \geq 0$ and $g-m \geq 1$, then by a result of Wiman [4, Theorem 6], the highest order of a cyclic action on $S_g(\mathcal{C})$ is $4(g-m) + 2$. If $r \geq 0$ and $g = m$, then consider the permuting $(n, 2r, 2k)$ -action t on S_0 that is nonseparating with respect to $\mathcal{C}$ guaranteed by Theorem 4.3. Since $t \in \text{Mod}(S_0)$ is an element of order n , it must be a rotation by $2\pi/n$ radians. Since the two fixed points of this action are not compatible in the sense of Definition 4.2, $r = 0$ and every non-trivial orbit has size n . Hence, $m = nk$ and so $n \mid m$, and in particular, $n \leq m = g$. Furthermore, if $m = g$, let t be the rotation of the sphere by $2\pi/n$, and $c \in \mathcal{C}$ be any curve. Then $h := tc$ is a root of $t_{\mathcal{C}}$ of degree m and (i) follows.

(ii) and (iii) follow from [7, Corollary 2.2]. If $r \geq 2$, then consider the corresponding permuting $(n, 2r, 2k)$ -data set from Theorem 3.8. Note that the genus $(g-m)$ of the permuting data set is given by

$$\frac{2 - 2(g-m)}{n} = 2 - 2g_0 + 2r \left(\frac{1}{n} - 1 \right) + \sum_{j=1}^s \left(\frac{1}{n_j} - 1 \right).$$

Since $(1 - 1/n_j) \geq 0$ and $g_0 \geq 0$, we have $(g-m) - 1 + r \geq n(-1 + r)$, from which (iv) follows. $\square$

5. SEPARATING MULTICURVES

A separating multicurve $\mathcal{C}$ in S_g is the disjoint union of finitely many pseudo-nonseparating curves. In this case $S_g(\mathcal{C})$ is disconnected, so we will require multiple finite order actions on the individual components of $S_g(\mathcal{C})$ to come together to form a root of $t_{\mathcal{C}}$ on S_g . To improve the exposition, we begin with the case $\mathcal{C} = \mathcal{C}^{(m)}$ so that $S_g(\mathcal{C})$ has two components.

Definition 5.1. Equivalence classes $\llbracket t_i \rrbracket$ of permuting (n_i, r_i, k_i) -actions on S_{g_i} , for $i = 1, 2$, are said to form an (r, k) -compatible pair $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ of degree n for integers $r, k \geq 0$, if

(i) $n = \text{lcm}(n_1, n_2)$, and

(ii) there exists $\{P_{i,1}, P_{i,2}, \dots, P_{i,r}\} \subset \mathbb{P}(t_i)$ such that for $1 \leq j \leq r$,

$$\theta_{P_{1,j}}(t_1) + \theta_{P_{2,j}}(t_2) = \frac{2\pi}{n} \pmod{2\pi}, \text{ and}$$

(iii) there exists $\{\mathbb{O}_{i,1}, \mathbb{O}_{i,2}, \dots, \mathbb{O}_{i,k}\} \subset \mathbb{O}(t_i)$ for $i = 1, 2$ such that $\mathbb{O}_{1,j} \sim \mathbb{O}_{2,j}$ as in Definition 4.1, for $1 \leq j \leq k$.

If

$$\alpha := \sum_{j=1}^k |\mathbb{O}_{1,j}| = \sum_{j=1}^k |\mathbb{O}_{2,j}|$$

then the number $g := g_1 + g_2 + r + \alpha - 1$ is called the *genus* of the pair $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$. Note that two actions are $(1,0)$ -compatible if they are compatible as nested actions in the sense of [9, Definition 3.2].

Notation 5.2. Let $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ be an (r, k) -compatible pair of degree n as in Definition 5.1. We write $\mathbb{P}(t_1, t_2) := \{P_{1,1}, P_{1,2}, \dots, P_{1,r}\}$ and $\mathbb{O}(t_1, t_2) := \{\mathbb{O}_{1,1}, \mathbb{O}_{1,2}, \dots, \mathbb{O}_{1,k}\}$. We define $\mathbb{P}(t_2, t_1)$ and $\mathbb{O}(t_2, t_1)$ similarly.

Lemma 5.3. *Let $\mathcal{C} = \{c_1, c_2, \dots, c_m\}$ be a multicurve on S_g , and N_i be annular neighbourhood of c_i . Write $N = \sqcup_{i=1}^m N_i$, and suppose $t \in \text{Mod}(S_g)$ is such that $t|_{S_g \setminus N} = \text{id}_{S_g \setminus N}$, then $\exists d_1, d_2, \dots, d_m \in \mathbb{N} \cup \{0\}$ such that $t = t_{c_1}^{d_1} \dots t_{c_m}^{d_m}$.*

Proof. Since $t|_{S_g \setminus N} = \text{id}_{S_g \setminus N}$, it follows that t fixes ∂N . The lemma now follows from the fact [1, Lemma 4.1.A, Chapter 12] that

$$\text{Mod}(N \text{fix}(\partial N)) \cong \oplus_{i=1}^m \text{Mod}(N_i \text{fix}(\partial N_i)) = \oplus_{i=1}^m \langle t_{c_i} \rangle.$$

□

Theorem 5.4. *Suppose that $S_g = S_{g_1} \#_{\mathcal{C}} S_{g_2}$, where $\mathcal{C} = \mathcal{C}^{(m)}$. Then (r, k) -permuting roots of $t_{\mathcal{C}}$ of degree n correspond to the (r, k) -compatible pairs $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ of equivalence classes of permuting (n_i, r_i, k_i) -actions on the S_{g_i} , of degree n .*

Proof. As before, we assume $m > 1$ and first show that every (r, k) -permuting root h of $t_{\mathcal{C}}$ of degree n yields a compatible pair $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ of degree n . Consider the (r, k) -partition $\mathcal{C}_{r,k}(h) = \{\mathcal{C}'_1, \dots, \mathcal{C}'_r, \mathcal{C}_1, \dots, \mathcal{C}_k\}$ of $\mathcal{C}$ induced by h . Let $\widehat{S_{g_s}}$ for $s = 1, 2$ denote the two components of $\widehat{S_g(\mathcal{C})}$. Let N be a closed annular neighborhood of $\mathcal{C}$. By isotopy, we may assume that $t_{\mathcal{C}}(\mathcal{C}) = \mathcal{C}$, $t_{\mathcal{C}}(N) = N$, and $t_{\mathcal{C}}|_{\widehat{S_{g_s}}} = \text{id}_{\widehat{S_{g_s}}}$. Putting $\widehat{t_s} = h|_{\widehat{S_{g_s}}}$, we may assume up to isotopy that $\widehat{t_s}^n|_{\widehat{S_{g_s}}} = \text{id}_{\widehat{S_{g_s}}}$ for $s = 1, 2$.

Let n_s be the smallest positive integer such that $\widehat{t_s}^{n_s} = \text{id}_{\widehat{S_{g_s}}}$ for $s = 1, 2$, and let $q = \text{lcm}(n_1, n_2)$. Then $t := h^q$ satisfies the hypotheses of Lemma 5.3. Hence, $\exists d_c \in \mathbb{N} \cup \{0\}$ such that

$$h^q = \prod_{c \in \mathcal{C}} t_c^{d_c}$$

Since $h^n|_{\widehat{S_{g_1}}} = \text{id}_{\widehat{S_{g_1}}}$ it follows that $n_1 \mid n$, and similarly $n_2 \mid n$. Hence, $q \mid n$ and so

$$\prod_{c \in \mathcal{C}} t_c = t_{\mathcal{C}} = (h^q)^{n/q} = \prod_{c \in \mathcal{C}} t_c^{nd_c/q}$$

Fix $c \in \mathcal{C}$ and restrict the functions on both sides of this equation to a closed annular neighbourhood of c disjoint from other curves in $\mathcal{C}$. As in Remark 2.3, we see that $nd_c/q = 1$ and hence $n = q = \text{lcm}(n_1, n_2)$.

We fill in $\partial \widehat{S_{g_s}}$ with disks to obtain the closed oriented surfaces S_{g_s} for $s = 1, 2$. We then extend $\widehat{t_s}$ to the S_{g_s} by coning. Thus t_s defines an effective C_{n_s} -action on the S_{g_s} , where $n_s \mid n$ for $s = 1, 2$, and $n = \text{lcm}(n_1, n_2)$.

When $r > 0$, the homeomorphism t_s fixes the center points $\{P_{s,1}, P_{s,2}, \dots, P_{s,r}\}$ of r disks in $S_{g_s} \setminus \widehat{S_{g_s}}$ for $s = 1, 2$. Hence we may write $\mathbb{P}(t_1, t_2) = \{P_{1,1}, P_{1,2}, \dots, P_{1,r}\}$ and $\mathbb{P}(t_2, t_1) = \{P_{2,1}, P_{2,2}, \dots, P_{2,r}\}$. For $1 \leq j \leq r$, the proof of [9, Theorem 3.4] implies that the corresponding turning angles around $P_{1,j}$ and $P_{2,j}$ must be compatible in the sense of condition (ii) of Definition 5.1.

When $k > 0$, let $\mathcal{C}_i = \{c_{i,1}, c_{i,2}, \dots, c_{i,m_i}\}$, for $1 \leq i \leq k$. Associated with each curve $c_{i,j} \in \mathcal{C}_i$, is a disk $D_{i,j}^s$ in each $S_{g_s} \setminus \widehat{S_{g_s}}$ for $s = 1, 2$. The centers $Q_{i,j}^s$ of the m_i disks $D_{i,j}^s$ for $1 \leq j \leq m_i$ form an orbit $\mathbb{O}_{s,i}$ in S_{g_s} for $s = 1, 2$. Thus we obtain a collection $\mathbb{O}_{t_s} = \{\mathbb{O}_{s,1}, \dots, \mathbb{O}_{s,k}\}$ of k distinguished nontrivial orbits on S_{g_s} for $s = i, j$. It remains to show that $\mathbb{O}_{1,i} \sim \mathbb{O}_{2,i}$, for $1 \leq i \leq k$, but the argument for this is similar to that of Theorem 4.3. Hence, the pair $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ forms an (r, k) -compatible pair of degree n .

The argument for the converse, and the fact that the resulting root is determined upto conjugacy, is analogous to that of Theorem 4.3. $\square$

Definition 5.5. Permuting data sets $\widetilde{\mathcal{D}}_i := (\mathcal{D}_i, \mathbb{O}_{\mathcal{D}_i})$ for $i = 1, 2$, where

$$\mathcal{D}_i = (n_i, \widetilde{g}_i, (a_{i,1}, \dots, a_{i,r_i}); ((c_{i,1}, y_{i,1}), \dots, (c_{i,s_i}, y_{i,s_i}))),$$

are said to be (r, k) -compatible of degree n for integers $r \geq 0$ and $k \geq 0$, if

- (i) $n = \text{lcm}(n_1, n_2)$,
- (ii) $r \leq r_i$, for $i = 1, 2$, and for $1 \leq j \leq r$,

$$a_{1,j} + a_{2,j} \equiv a_{1,j} a_{2,j} \pmod{n},$$

- (iii) there exists $S_i \subseteq \mathbb{O}_{\mathcal{D}_i}$ such that

(a)

$$k = \sum_{(p, m_p) \in S_1} m_p = \sum_{(q, m_q) \in S_2} m_q,$$

- (b) for every $(p, m_p) \in S_1$, there exists $(q, m_q) \in S_2$ such that

$$\theta(p) + \theta(q) \equiv \begin{cases} 0 & : \text{ if } p = q = (0, 1) \\ 2\pi/n \pmod{2\pi} & : \text{ otherwise} \end{cases}$$

If

$$\alpha := \sum_{((a,b), m_p) \in S_1} n_1/b = \sum_{((c,d), m_q) \in S_2} n_2/d$$

and if g_i is the genus of $\mathcal{D}_i$, then the number $g = g_1 + g_2 + r + \alpha - 1$ is called *genus* of the pair $(\widetilde{\mathcal{D}}_1, \widetilde{\mathcal{D}}_2)$.

The next theorem and its corollary follow directly from Theorems 3.8 and 5.4.

Theorem 5.6. *Suppose that $S_g = S_{g_1} \#_{\mathcal{C}} S_{g_2}$, where $\mathcal{C} = \mathcal{C}^{(m)}$. Then (r, k) -compatible pairs $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ of equivalence classes of permuting (n_i, r_i, k_i) -actions on the S_{g_i} correspond to (r, k) -compatible pairs $(\widetilde{\mathcal{D}}_1, \widetilde{\mathcal{D}}_2)$ of data sets of genus g , where $\widetilde{\mathcal{D}}_i$ is a permuting (n_i, r_i, k_i) -data set of genus g_i .*

Corollary 5.7. *Suppose that $S_g = S_{g_1} \#_{\mathcal{C}} S_{g_2}$, where $\mathcal{C} = \mathcal{C}^{(m)}$. Then conjugacy classes of (r, k) -permuting roots of $t_{\mathcal{C}}$ of degree n correspond to (r, k) -compatible pairs $(\widetilde{\mathcal{D}}_1, \widetilde{\mathcal{D}}_2)$ of degree n and genus g , where $\widetilde{\mathcal{D}}_i$ is a permuting (n_i, r_i, k_i) data set of genus g_i .*

We now consider the case of a $(0, k')$ -permuting root h of $t_{\mathcal{C}}$ where $\mathcal{C} = \mathcal{C}^{(k)}(m)$. Here, the restriction of h to $S_g(\mathcal{C})$ induces a non-trivial permutation of the components of $S_g(\mathcal{C})$. Since h is a homeomorphism, it maps one component to another of the same genus, thus inducing an action on each subsurface of the form $\mathbb{S}_g(m) \subset S_g(\mathcal{C})$. Therefore, we generalize the notion of a permuting (n, r, k) -action to encompass the action on $\mathbb{S}_g(m)$ induced by h .

Definition 5.8. Fix integers $g \geq 0$ and $m \geq 1$.

- (i) A homeomorphism $\sigma_m : \mathbb{S}_g(m) \rightarrow \mathbb{S}_g(m)$ is said to be *essential* if for all i , $\sigma_m|_{S_g^i}(S_g^i) = S_g^{i+1}$. An essential homeomorphism σ_m on $\mathbb{S}_g(m)$ can be viewed simply as the m -cycle $(1, 2, \dots, m)$ permuting its m components S_g^i .
- (ii) An orientation-preserving C_n -action t on $\mathbb{S}_g(m)$ is said to be a *permuting (n, r, k) -action* if $m \mid n$ and there exists an essential homeomorphism $\sigma_m : \mathbb{S}_g(m) \rightarrow \mathbb{S}_g(m)$ such that $t = \sigma_m \circ \tilde{t}$, where $\tilde{t}$ is a permuting (n, r, k) -action on each S_g^i .

Remark 5.9. Suppose that $\tilde{t} \in \text{Mod}(S_g)$ defined a permuting (n, r, k) -action and $t = \sigma_m \circ \tilde{t}$, then $t_i := t^m|_{S_g^i} \in \text{Mod}(S_g^i)$ defines a permuting $(n/m, \tilde{r}, \tilde{k})$ -action on S_g^i . Furthermore, all the t_i are conjugate to each other via σ_m .

Conversely, if $t' \in \text{Mod}(S_g)$ is a permuting $(n/m, \tilde{r}, \tilde{k})$ -action on S_g which has an m^{th} root $\tilde{t} \in \text{Mod}(S_g)$, then the map $t := \sigma_m \circ \tilde{t}$ defines a permuting (n, r, k) -action on $\mathbb{S}_g(m)$. Thus, a permuting (n, r, k) -action on $\mathbb{S}_g(m)$ corresponds to a permuting $(n/m, \tilde{r}, \tilde{k})$ -action on S_g which has an m^{th} root in $\text{Mod}(S_g)$.

Definition 5.10. Let t_1 and t_2 be two permuting (n, r, k) -actions on $\mathbb{S}_g(m)$. Then we say t_1 is *equivalent* to t_2 if $t_1^m|_{S_g^i}$ and $t_2^m|_{S_g^i}$ are equivalent as permuting $(n/m, \tilde{r}, \tilde{k})$ -actions on S_g^i in the sense of Definition 3.4.

Definition 5.11. Let $u, v \geq 0$ be fixed integers, t_1 be a permuting (n_1, r_1, k_1) -action on S_{g_1} and t_2 be a permuting (n_2, r_2, k_2) -action on $\mathbb{S}_{g_2}(m)$ such that

$t_2 = \sigma_m \circ \tilde{t}_2$ as in Remark 5.9. Then the equivalence classes $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ are said to form a (u, v) -compatible pair if

- (i) for $1 \leq i \leq m$, $(\llbracket t_1^m \rrbracket, \llbracket t_2^m|_{S_{g_2}^i} \rrbracket)$ is an (u, v) -compatible pair of degree n/m , where $n = \text{lcm}(n_1, n_2)$,
- (ii) for $1 \leq i \leq m, 1 \leq j \leq u$, there mutually disjoint pairs $\{P_{i,j}^1, P_{i,j}^2\}$, where $P_{i,j}^1 \in \mathbb{P}(t_1^m)$ and $P_{i,j}^2 \in \mathbb{P}(t_2^m|_{S_{g_2}^i})$ such that

$$\theta_{P_{i,j}^1}(t_1) + \theta_{P_{i,j}^2}(\tilde{t}_2) = 2\pi/n \pmod{2\pi}, \text{ and}$$

- (iii) for $1 \leq i \leq m, 1 \leq j \leq v$, mutually disjoint pairs $\{\mathbb{O}_{i,j}^1, \mathbb{O}_{i,j}^2\}$, where $\mathbb{O}_{i,j}^1 \in \mathbb{O}(t_1)$ and $\mathbb{O}_{i,j}^2 \in \mathbb{O}(\tilde{t}_2)$, such that $\mathbb{O}_{i,j}^1 \sim \mathbb{O}_{i,j}^2$, as in Definition 4.1.

The number $n = \text{lcm}(n_1, n_2)$ is called the *degree* of the pair, and the number $g = g_1 + m(g_2 + k - 1)$ is called the *genus* of the pair.

Notation 5.12. Let $(\llbracket t_1 \rrbracket, \llbracket t_2 \rrbracket)$ be a (u, v) -compatible pair of degree n as in Definition 5.11. We write

- (i) $\mathbb{P}(t_1, t_2) = \{P_{i,j}^1 : 1 \leq i \leq m, 1 \leq j \leq u\}$ and $\mathbb{P}(t_2, t_1) = \{P_{i,j}^2 : 1 \leq i \leq m, 1 \leq j \leq u\}$.
- (ii) Similarly, we define $\mathbb{O}(t_1, t_2) = \{\mathbb{O}_{i,j}^1 : 1 \leq i \leq m, 1 \leq j \leq v\}$ and $\mathbb{O}(t_1, t_2) = \{\mathbb{O}_{i,j}^2 : 1 \leq i \leq m, 1 \leq j \leq v\}$.

Lemma 5.13. Suppose that $\mathcal{C} = \mathcal{C}^{(k)}(m)$ be a multicurve in S_g . Then conjugacy classes of $(0, k')$ -permuting roots of $t_{\mathcal{C}}$ of degree n correspond to (u, v) -compatible pairs of degree n and genus g , where $k' = u + v$.

Proof. Let $h \in \text{Mod}(S_g)$ be a $(0, k')$ -permuting root of degree n on S_g . Then as in Theorem 5.4, we obtain an effective C_{n_1} -action t_1 on S_{g_1} and an effective C_{n_2} -action on $S_{g_2}(m)$, where $n = \text{lcm}(n_1, n_2)$. Furthermore, h restricts to an essential homeomorphism $\sigma_m : \mathbb{S}_g(m) \rightarrow \mathbb{S}_g(m)$ such that $\sigma_m \circ t_2 = t_2 \circ \sigma_m$. Hence the maps

$$t_{2,i} := \sigma_m^{-1} t|_{S_{g_2}^i} : S_{g_2}^i \rightarrow S_{g_2}^i$$

are conjugate to each other when considered as elements of $\text{Mod}(S_g)$, and so t_2 is a permuting (n_2, r_2, k_2) -action on $S_{g_2}(m)$ as in Definition 5.8. Since h^m is a root of $t_{\mathcal{C}}$ of degree (n/m) that preserves each $\mathcal{C}_i^{(k)}$, we may restrict h^m and cone to obtain maps

$$h_i \in \text{Mod}(\Sigma_i), \text{ where } \Sigma_i := S_{g_1} \#_{\mathcal{C}_i^{(k)}} S_{g_2}^i,$$

which are pairwise conjugate to each other via h . Thus, it follows from Theorem 5.4 that there exist integers $u, v \geq 0$ such that $(\llbracket t_1^m \rrbracket, \llbracket t_2^m|_{S_{g_2}^i} \rrbracket)$ forms a (u, v) -compatible pair of degree n/m , for $1 \leq i \leq m$.

By condition (ii) of Definition 5.1, there exists mutually disjoint pairs $\{P_{i,j}^1, P_{i,j}^2\}$, where $P_{i,j}^1 \in \mathbb{P}(t_1^m)$ and $P_{i,j}^2 \in \mathbb{P}(t_2^m|_{S_{g_2}^i})$, such that

$$\theta_{P_{i,j}^1}(t_1^m) + \theta_{P_{i,j}^2}(\tilde{t}_2^m) = \frac{2\pi}{(n/m)} \pmod{2\pi}.$$

Once again, since h^m is a root of $t_{\mathcal{C}}$ it follows that

$$\theta_{P_{i,j}^1}(t_1) + \theta_{P_{i,j}^2}(\tilde{t}_2) = 2\pi/n \pmod{2\pi}.$$

Similarly, one obtains condition (iii) of Definition 5.11 as well. Note that every fixed point of h^m , and every orbit of h^m , induces one orbit of h , and thus $k' = u + v$. The converse is a just a matter of reversing this argument. $\square$

We now consider the case of an (r, k) -permuting root of $t_{\mathcal{C}}$ where $r, k > 0$. We begin by writing S_g as a connected sum of subsurfaces S_{g_i} across those pseudo-nonseparating submulticurves which are preserved by h . The restriction of h to each S_{g_i} is then a $(0, k_i)$ -permuting root of the Dehn twist about the submulticurve $\mathcal{C} \cap S_{g_i}$. This allows us to apply Theorem 5.13 to obtain finite order actions on S_{g_i} such that pairs of actions on adjacent subsurfaces are compatible (in the sense of Theorem 5.4).

Notation 5.14. Let $\mathcal{C}$ be a separating multicurve in S_g , and let h be an (r, k) -permuting root of $t_{\mathcal{C}}$.

- (i) We shall denote the set of all pseudo-nonseparating submulticurves of $\mathcal{C}$ that are preserved by h by $\text{Fix}_h(\mathcal{C}) = \{\mathcal{E}_1, \dots, \mathcal{E}_{m(h)}\}$.
- (ii) Writing

$$S_g = \overline{\#}_{i=1}^{m(h)} (S_{g_i} \#_{\mathcal{E}_i} S_{g_{i+1}})$$

and $D_i := \mathcal{C} \cap S_{g_i}$, we have that for each i , $S_{g_i} \cap \mathcal{C}_{r,k}(h)$ is a $(0, k_i)$ -partition of D_i , which has the form

$$S_{g_i} \cap \mathcal{C}_{r,k}(h) = \{\mathcal{C}_{i,j}^{(k_{i,j})}(m_{i,j}) : 1 \leq j \leq k_i\}.$$

- (iii) For $1 \leq i \leq r+1$, we write

$$S_{g_i} = \overline{\#}_{j=1}^{k_i} (S_{g_{i,1}} \#_{D_{i,j}} S_{g_{i,2,j}}(m_{i,j})),$$

where $D_{i,j} = \mathcal{C}_{i,j}^{(k_{i,j})}(m_{i,j})$ and $g_i = g_{i,1} + \sum_{j=1}^{k_i} m_{i,j}(g_{i,2,j} + k_{i,j} - 1)$.


- (iv) We will denote $\min_{1 \leq i \leq m(h)} g_i + g_{i+1}$ by $g(\mathcal{C})$.

In Figure 6 below, we have the surface S_{24} with the multicurve

$$\mathcal{C} = \mathcal{C}^{(2)}(2) \sqcup \mathcal{C}^{(1)}(2) \sqcup \mathcal{C}^{(3)} \sqcup \mathcal{C}^{(1)} \sqcup \mathcal{C}^{(1)}(3).$$

According to Notation 5.14,

$$S_{24} = \left(\mathbb{S}_3(2) \#_{\mathcal{C}^{(2)}(2)} S_2 \#_{\mathcal{C}^{(1)}(2)} \mathbb{S}_1(2) \right) \#_{\mathcal{C}^{(3)}} S_2 \#_{\mathcal{C}^{(1)}} \left(S_1 \#_{\mathcal{C}^{(1)}(3)} \mathbb{S}_3(3) \right).$$

FIGURE 6. S_{24} with a separating multicurve $\mathcal{C}$.

Definition 5.15. Let t_1 be a permuting $(n_1, 0, k_1)$ -action on S_{g_1} , and let $t_{2,j}$ be a permuting $(n_{2,j}, 0, k_{2,j})$ -action on $\mathbb{S}_{g_{2,j}}(m_j)$, for $1 \leq j \leq s$. Then $(\llbracket t_1 \rrbracket, \llbracket t_{2,1} \rrbracket, \dots, \llbracket t_{2,s} \rrbracket)$ forms a $(s+1)$ -compatible tuple of degree n if

- (i) for each $1 \leq j \leq s$, $(\llbracket t_1 \rrbracket, \llbracket t_{2,j} \rrbracket)$ forms an (u_j, v_j) -compatible pair of degree n , for some $u_j, v_j \geq 0$ such that $k_{2,j} = u_j + v_j$.
- (ii) For each $i \neq j$,

$$\mathbb{O}(t_1, t_{2,i}) \cap \mathbb{O}(t_1, t_{2,j}) = \emptyset = \mathbb{P}(t_1, t_{2,i}) \cap \mathbb{P}(t_1, t_{2,j}).$$

The number $g = g_1 + \sum_{j=1}^s m_j(g_{2,j} + k_{2,j} - 1)$ is called the *genus* of the $(s+1)$ -tuple.

Definition 5.16. Fix $m, n \in \mathbb{N}$, and for $1 \leq i \leq m+1$, let

$$\bar{t}_i = (\llbracket t_{i,1} \rrbracket, \llbracket t_{i,2,1} \rrbracket, \dots, \llbracket t_{i,2,s_i} \rrbracket)$$

be a $(s_i + 1)$ -compatible tuple as in Definition 5.15. Then the tuple

$$(\bar{t}_1, \bar{t}_2, \dots, \bar{t}_{m+1})$$

is said to form an $(m+1)$ -compatible multituple of degree n if for each $1 \leq i \leq (m+1)$,

- (i) the pair $(\llbracket t_{i,1} \rrbracket, \llbracket t_{i+1,1} \rrbracket)$ forms an $(r_{i,1}, k_{i,1})$ -compatible pair of degree n , and
- (ii) $\mathbb{O}(t_{i,1}, t_{i+1,1}) \cap \left(\sqcup_{j=1}^{s_i} \mathbb{O}(t_{i,1}, t_{i,2,j}) \right) = \emptyset = \mathbb{P}(t_{i,1}, t_{i+1,1}) \cap \left(\sqcup_{j=1}^{s_i} \mathbb{P}(t_{i,1}, t_{i,2,j}) \right)$.

If $g(\bar{t}_i)$ denotes the genus of $\bar{t}_i$, and

$$\alpha_i := \sum_{\mathbb{O} \in \mathbb{O}(t_{1,i}, t_{1,i+1})} |\mathbb{O}|$$

then the number

$$g = \sum_{i=1}^{m+1} g(\bar{t}_i) + \sum_{i=1}^m (r_{1,i} + k_{1,i} \alpha_i - 1)$$

is called the *genus* of the multituple.

The following theorem follows from Lemmas 5.4 and 5.13.

Theorem 5.17. Let $\mathcal{C}$ be separating multicurve in S_g . Then the conjugacy class of a root h of $t_{\mathcal{C}}$ of degree n with $|\text{Fix}_h(\mathcal{C})| = m$ corresponds to an $(m+1)$ -compatible multituple of degree n and genus g .

The following corollary, which gives an upper bound for the degree of a root of t_C , follows from [9, Theorem 8.6] and Theorem 5.17.

Corollary 5.18. *Let C be a separating multicurve in S_g and h be a root of t_C of degree n . Then $n \leq 4g(C)^2 + 2g(C)$.*

The following corollary, which gives a stable upper bound for the degree of a root of t_C , follows from [9, Theorem 8.14] and Theorem 5.17.

Corollary 5.19. *Let C be a separating multicurve in S_g and h be a root of t_C of degree n . Then $n \leq \frac{16}{5}g^2 + 12g + \frac{45}{4}$, if $g(C) \geq 10$.*

The primary purpose behind developing the notion of data sets was for algebraically describing actions that are involved in the construction of roots. Owing to complexity of notation, we shall abstain from developing separate notation involving data sets for encoding compatible tuples or multituples. We will see later in Sections 6 and 7 that one can easily circumvent the need for complex notation by a judicious use of Remark 5.9.

Recall that a multicurve C is said to be mixed if it is neither separating nor nonseparating. For the sake of brevity, we do not develop a separate theory to describe roots of Dehn twists about mixed multicurves as such a theory would merely be a combination of the separating and nonseparating cases. However, we will classify such roots on S_3 and S_4 , thereby indicating how such a theory would follow from the ideas developed in Sections 4 and 5.

6. CLASSIFICATION OF ROOTS FOR S_3

In this, and the subsequent section, we classify roots of Dehn twists about multicurves on surfaces of genus 3 and 4. Once again, we break up the classification into nonseparating, separating and mixed multicurves.

When classifying an $(m+1)$ -compatible multituples $(\overline{t_1}, \overline{t_2}, \dots, \overline{t_{m+1}})$ that corresponds to a root, Condition (i) of Definition 5.16 and Condition (ii) of Definition 5.15 help in eliminating data sets that do not lead to roots. For the sake of brevity, we only list those data sets that do lead to roots. Furthermore, in each case, a careful examination of the data set $\mathcal{D}$ also gives $\mathbb{O}_{\mathcal{D}}$, and so we only display the former.

Finally, when $\overline{t_i}$ is a permuting (n, r, k) -action on $\mathbb{S}_g(m)$, we use Remark 5.9 and replace $\overline{t_i}$ by the corresponding action on S_g which has an m^{th} root $\tilde{t_i}$, whose equivalence class can be encoded by a data set D_i . Therefore an $(m+1)$ -compatible multituple is described by a tuple $(D_1, D_2, \dots, D_{m+1})$ of data sets, which will be listed in a table. While enumerating the curves in a multicurve, we use the convention that separating curves are denoted with the letter c , while nonseparating curves are denoted with the letter d .

C is a nonseparating multicurve.

Degree	D_1
3	$(3, 0; (1, 3), (1, 3))$

Table 1: $\mathcal{C} = \{d_1, d_2, d_3\}$ and $(r, k) = (0, 1)$

Degree	D_1
2	$(2, 1;)$

Table 2: $\mathcal{C} = \{d_1, d_2\}$ and $(r, k) = (0, 1)$

$\mathcal{C}$ is a separating multicurve.

Degree	D_1	D_2
3	$(3, 0; (1, 3), (1, 3))$	$(1, 1;)$

Table 3: $\mathcal{C} = \{c_1, c_2, c_3\}$, $S_3 = S_0 \#_{\mathcal{C}} S_1(3)$ and $(r, k) = (0, 1)$

Degree	D_1	D_2	D_3
2	$(1, 1, 1;)$	$(2, 0, 1; (1, 2))$	$(1, 1;)$

Table 4: $\mathcal{C} = \{c_1, c_2, c_3\}$, $S_3 = S_0 \#_{c_1} S_1 \#_{\{c_2, c_3\}} S_1(2)$ and $(r, k) = (1, 1)$

Degree	D_1	D_2	D_3
3	$(3, 0, 2; (2, 3), (2, 3))$	$(3, 0, (2, 2); (2, 3))$	$(3, 0, 2; (2, 3), (2, 3))$
6	$(6, 0, 5; (1, 2), (2, 3))$	$(3, 0, (1, 1); (1, 3))$	$(6, 0, 5; (1, 2), (2, 3))$
6	$(6, 0, 1; (1, 2), (1, 3))$	$(1, 1, (1, 1);)$	$(6, 0, 1; (1, 2), (1, 3))$
6	$(3, 0, 2; (2, 3), (2, 3))$	$(2, 0, (1, 1); (1, 2), (1, 2))$	$(3, 0, 2; (2, 3), (2, 3))$
6	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$	$(3, 0, (2, 2); (2, 3))$	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$
12	$(3, 0, 1; (1, 3), (1, 3))$	$(4, 0, (3, 3); (1, 2))$	$(3, 0, 1; (1, 3), (1, 3))$
12	$(4, 0, 3; (1, 2), (3, 4))$	$(3, 0, (1, 1); (1, 3))$	$(4, 0, 3; (1, 2), (3, 4))$

Table 5: $\mathcal{C} = \{c_1, c_2\}$, $S_3 = S_1 \#_{c_1} S_1 \#_{c_2} S_1$ and $(r, k) = (2, 0)$

Degree	D_1	D_2
2	$(2, 1;)$	$(1, 1;)$
12	$(4, 0; (1, 4), (1, 2), (1, 4))$	$(6, 0; (5, 6), (1, 2), (2, 3))$
12	$(4, 0; (3, 4), (1, 2), (3, 4))$	$(6, 0; (5, 6), (1, 2), (2, 3))$
12	$(6, 0; (5, 6), (1, 2), (2, 3))$	$(4, 0; (1, 4), (1, 2), (1, 4))$
12	$(6, 0; (5, 6), (1, 2), (2, 3))$	$(4, 0; (3, 4), (1, 2), (3, 4))$

Table 6: $\mathcal{C} = \{c_1, c_2\}$, $S_3 = S_1 \#_{\mathcal{C}} S_1(2)$ and $(r, k) = (0, 1)$

Degree	D_1	D_2
2	$(2, 0, (1, 1); (1, 2), (1, 2), (1, 2))$	$(1, 0, (1, 1);)$
3	$(3, 0, (1, 1); (1, 3))$	$(1, 0, (1, 1);)$
3	$(3, 0, (2, 2); (2, 3))$	$(3, 0, (2, 2); (2, 3))$
4	$(4, 0, (1, 1); (1, 2))$	$(1, 0, (1, 1);)$
4	$(4, 0, (3, 3); (1, 2))$	$(2, 0, (1, 1); (1, 2), (1, 2), (1, 2))$
6	$(3, 0, (2, 2); (2, 3))$	$(2, 0, (1, 1); (1, 2), (1, 2), (1, 2))$
12	$(4, 0, (3, 3); (1, 2))$	$(3, 0, (1, 1); (1, 3))$

Table 7: $\mathcal{C} = \mathcal{C}^{(2)}$, $S_3 = S_1 \#_{\mathcal{C}} S_1$, and $(r, k) = (2, 0)$

Degree	D_1	D_2
6	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2))$	$(6, 0; (1, 6), (1, 2), (1, 3))$

Table 8: $\mathcal{C} = \mathcal{C}^{(2)}$, $S_3 = S_1 \#_{\mathcal{C}} S_1$ and $(r, k) = (0, 1)$

$\mathcal{C}$ is a mixed multicurve.

Degree	D_1	D_2
2	$(3, 0; (1, 3), (1, 3))$	$(1, 1;)$

Table 9: $\mathcal{C} = \{c_1, c_2, c_3, d_1, d_2, d_3\}$, $S_3 = S_0 \#_{\{c_1, c_2, c_3\}} \# S_1(3)$ where $d_i \in \widehat{S}_1^i$ and $(r, k) = (0, 2)$

Degree	D_1	D_2	D_3
2	$(1, 1, 1;)$	$(2, 0, 1; (1, 2))$	$(1, 1;)$

Table 10: $\mathcal{C} = \{c_1, c_2, c_3, d_1, d_2\}$, $S_3 = S_1 \#_{c_1} S_0 \#_{\{c_2, c_3\}} S_1(2)$ where $d_i \in \widehat{S}_1^i$ for $i = 1, 2$ and $(r, k) = (1, 2)$

Degree	D_1	D_2
2	$(2, 1;)$	$(2, 0;)$

Table 11: $\mathcal{C} = \{c_1, c_2, d_1, d_2\}$, $S_3 = S_1 \#_{\{c_1, c_2\}} S_1(2)$ and $d_i \in \widehat{S}_1^i$ for $i = 1, 2$ and $(r, k) = (0, 2)$

Degree	D_1	D_2
2	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$	$(1, 1, 1;)$

Table 12: $\mathcal{C} = \{c_1, d_1, d_2\}$, $S_3 = S_1 \#_{c_1} S_2$ and $\{d_1, d_2\} \subset \widehat{S}_2$ and $(r, k) = (1, 1)$

Degree	D_1	D_2
3	$(3, 0, 2; (2, 3), (2, 3))$	$(3, 0, (2, 2, 2);)$

Table 13: $\mathcal{C} = \{c_1, d_1\}$ mixed and $S_3 = S_1 \#_{c_1} S_2$ where $d_1 \in \widehat{S_2}$ and $(r, k) = (2, 0)$

7. CLASSIFICATION OF ROOTS FOR S_4

$\mathcal{C}$ is a nonseparating multicurve.

Degree	D_1
4	$(4, 0; (1, 4), (1, 4))$
2	$(2, 1; (1, 2), (1, 2))$

Table 14: $\mathcal{C} = \{d_1, d_2, d_3, d_4\}$ and $(r, k) = (0, 1)$

Degree	D_1
3	$(3, 1;)$

Table 15: $\mathcal{C} = \{d_1, d_2, d_3\}$ and $(r, k) = (0, 1)$

Degree	D_1
2	$(2, 1; (1, 2), (1, 2))$

Table 16: $\mathcal{C} = \{d_1, d_2\}$ and $(r, k) = (0, 1)$

$\mathcal{C}$ is a separating multicurve.

Degree	D_1	D_2
4	$(4, 0; (1, 4), (1, 4))$	$(1, 1;)$

Table 17: $\mathcal{C} = \{c_1, c_2, c_3, c_4\}$, $S_4 = S_0 \#_{\mathcal{C}} S_1(4)$ and $(r, k) = (0, 1)$

Degree	D_1	D_2	D_3
2	$(1, 1;)$	$(2, 0;)$	$(1, 1;)$
6	$(6, 0; (1, 6), (1, 2), (1, 3))$	$(2, 0;)$	$(6, 0; (1, 6), (1, 2), (1, 3))$

Table 18: $\mathcal{C} = \{c_1, c_2, c_3, c_4\}$, $S_4 = S_1(2) \#_{\{c_1, c_2\}} S_0 \#_{\{c_3, c_4\}} S_1(2)$ and $(r, k) = (0, 2)$

Degree	D_1	D_2
3	(3, 1;)	(1, 1;)
6	(6, 0; (1, 6), (1, 2), (1, 3))	(1, 1;)
6	(6, 0; (5, 6), (1, 2), (2, 3))	(1, 1;)
6	(3, 1;)	(6, 0; (1, 6), (1, 2), (1, 3))
6	(3, 1;)	(6, 0; (5, 6), (1, 2), (2, 3))

Table 19: $\mathcal{C} = \{c_1, c_2, c_3\}$, $S_4 = S_1 \#_{\mathcal{C}} S_1(3)$ and $(r, k) = (0, 1)$.

Degree	D_1	D_2	D_3
4	(1, 1, 1;)	(4, 0, 1; (1, 2), (1, 4))	(1, 1;)
6	(1, 1, 1;)	(6, 0, 1; (1, 2), (1, 3))	(1, 1;)
6	(3, 0, 2; (2, 3), (2, 3))	(2, 0, 1; (1, 2), (1, 2), (1, 2))	(6, 0; (1, 6), (1, 2), (1, 3))

Table 20: $\mathcal{C} = \{c_1, c_2, c_3\}$, $S_4 = S_1 \#_{c_1} S_1 \#_{\{c_2, c_3\}} S_1(2)$ and $(r, k) = (1, 1)$

Degree	D1	D2	D3	D4
3	(3, 0, 2; (2, 3), (2, 3))	(3, 0, 2; (2, 3), (2, 3))	(3, 0, 2; (2, 3), (2, 3))	(3, 0, 2; (2, 3), (2, 3))
6	(6, 0, 5; (1, 2), (2, 3))	(3, 0, 1; (1, 3), (1, 3))	(6, 0, 5; (1, 2), (2, 3))	(3, 0, 1; (1, 3), (1, 3))
12	(3, 0, 1; (1, 3), (1, 3))	(4, 0, 3; (1, 2), (3, 4))	(3, 0, 1; (1, 3), (1, 3))	(4, 0, 3; (1, 2), (3, 4))
12	(4, 0, 3; (1, 2), (3, 4))	(3, 0, 1; (1, 3), (1, 3))	(4, 0, 3; (1, 2), (3, 4))	(3, 0, 1; (1, 3), (1, 3))

Table 21: $\mathcal{C} = \{c_1, c_2, c_3\}$, $S_4 = S_1 \#_{c_1} S_1 \#_{c_2} S_1 \#_{c_3} S_1$ and $(r, k) = (3, 0)$

Degree	D_1	D_2
6	(2, 0; (1, 2), (1, 2), (1, 2), (1, 2), (1, 2), (1, 2))	(6, 0; (1, 6), (1, 2), (1, 3))
6	(2, 1; (1, 2), (1, 2))	(6, 0; (1, 6), (1, 2), (1, 3))
6	(6, 0; (5, 6), (1, 3), (5, 6))	(1, 1;)

Table 22: $\mathcal{C} = \{c_1, c_2\}$, $S_4 = S_2 \#_{\mathcal{C}} S_1(2)$ and $(r, k) = (0, 1)$.

Degree	D_1	D_2	D_3
6	(6, 0, 5; (1, 2), (2, 3))	(3, 0, 1; (1, 3), (1, 3))	(6, 0, 5; (1, 3), (5, 6))

Table 23: $\mathcal{C} = \{c_1, c_2\}$, $S_4 = S_1 \#_{c_1} S_1 \#_{c_2} S_2$ and $(r, k) = (2, 0)$.

Degree	D_1	D_2
3	$(3, 0; (1, 3), (1, 3), (1, 3))$	$(3, 0; (1, 3), (1, 3), (1, 3))$
3	$(3, 0; (1, 3), (1, 3), (1, 3))$	$(3, 0; (2, 3), (2, 3), (2, 3))$
3	$(3, 0; (2, 3), (2, 3), (2, 3))$	$(3, 0; (2, 3), (2, 3), (2, 3))$
3	$(3, 0; (2, 3), (2, 3), (2, 3))$	$(3, 0; (1, 3), (1, 3), (1, 3))$
6	$(3, 0; (1, 3), (1, 3), (1, 3))$	$(6, 0; (1, 6), (1, 2), (1, 3))$
6	$(3, 0; (1, 3), (1, 3), (1, 3))$	$(6, 0; (5, 6), (1, 2), (2, 3))$
6	$(3, 0; (2, 3), (2, 3), (2, 3))$	$(6, 0; (1, 6), (1, 2), (1, 3))$
6	$(3, 0; (2, 3), (2, 3), (2, 3))$	$(6, 0; (5, 6), (1, 2), (2, 3))$

Table 24: $\mathcal{C} = \mathcal{C}^{(3)}$, $S_4 = S_1 \#_{\mathcal{C}} S_1$ and $(r, k) = (0, 1)$

Degree	D_1	D_2
4	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$	$(4, 0, 3; (1, 2), (3, 4))$

Table 25: $\mathcal{C} = \mathcal{C}^{(3)}$, $S_4 = S_1 \#_{\mathcal{C}} S_1$ and $(r, k) = (1, 1)$

Degree	D_1	D_2
3	$(3, 0, (2, 2, 2);)$	$(3, 0, (2, 2, 2);)$
6	$(3, 0, (2, 2, 2);)$	$(2, 0, (1, 1, 1); (1, 2))$

Table 26: $\mathcal{C} = \mathcal{C}^{(3)}$, $S_4 = S_1 \#_{\mathcal{C}} S_1$ and $(r, k) = (3, 0)$.

Degree	D_1	D_2	D_3
4	$(4, 0; (1, 4), (1, 2), (1, 4))$	$(2, 0; (1, 2), (1, 2))$	$(4, 0; (1, 4), (1, 2), (1, 4))$
4	$(4, 0; (1, 4), (1, 2), (1, 4))$	$(2, 0; (1, 2), (1, 2))$	$(4, 0; (3, 4), (1, 2), (3, 4))$
4	$(4, 0; (3, 4), (1, 2), (3, 4))$	$(2, 0; (1, 2), (1, 2))$	$(4, 0; (3, 4), (1, 2), (3, 4))$
6	$(6, 0; (1, 6), (1, 2), (1, 3))$	$(2, 0; (1, 2), (1, 2))$	$(6, 0; (1, 6), (1, 2), (1, 3))$
6	$(6, 0; (1, 6), (1, 2), (1, 3))$	$(2, 0; (1, 2), (1, 2))$	$(6, 0; (5, 6), (1, 2), (2, 3))$
6	$(6, 0; (5, 6), (1, 2), (2, 3))$	$(2, 0; (1, 2), (1, 2))$	$(6, 0; (5, 6), (1, 2), (2, 3))$

Table 27: $\mathcal{C} = \mathcal{C}^{(2)} \sqcup \mathcal{C}^{(2)}$, $S_4 = S_1 \#_{\mathcal{C}^{(2)}} S_0 \#_{\mathcal{C}^{(2)}} S_1$ and $(r, k) = (0, 2)$.

Degree	D_1	D_2
2	$(2, 0; (1, 2), (1, 2))$	$(1, 1;)$
6	$(2, 0; (1, 2), (1, 2))$	$(6, 0; (1, 6), (1, 2), (1, 3))$

Table 28: $\mathcal{C} = \mathcal{C}^{(2)} \sqcup \mathcal{C}^{(2)}$, $S_4 = S_0 \#_{\mathcal{C}} S_1(2)$ and $(r, k) = (0, 2)$.

Degree	D_1	D_2
4	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2), (1, 2), (1, 2))$	$(4, 0; (1, 4), (1, 2), (1, 4))$
4	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2), (1, 2), (1, 2))$	$(4, 0; (3, 4), (1, 2), (3, 4))$
4	$(2, 1; (1, 2), (1, 2))$	$(4, 0; (1, 4), (1, 2), (1, 4))$
4	$(2, 1; (1, 2), (1, 2))$	$(4, 0; (3, 4), (1, 2), (3, 4))$
4	$(4, 0; (1, 2), (1, 2), (3, 4))$	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2))$
4	$(4, 0; (1, 2), (1, 2), (3, 4))$	$(2, 1;)$
4	$(4, 0; (1, 2), (1, 2), (2, 4))$	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2))$
4	$(4, 0; (1, 2), (1, 2), (2, 4))$	$(2, 1;)$
6	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2), (1, 2), (1, 2))$	$(6, 0; (1, 6), (1, 2), (1, 3))$
6	$(2, 1; (1, 2), (1, 2))$	$(6, 0; (1, 6), (1, 2), (1, 3))$
6	$(6, 0; (5, 6), (1, 3), (5, 6))$	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2))$
6	$(6, 0; (5, 6), (1, 3), (5, 6))$	$(2, 1;)$

Table 29: $\mathcal{C} = \mathcal{C}^{(2)}$, $S_4 = S_2 \#_{\mathcal{C}} S_1$ and $(r, k) = (0, 1)$.

Degree	D_1	D_2
2	$(1, 2, (1, 1);)$	$(2, 0, (1, 1); (1, 2), (1, 2))$
2	$(2, 0, (1, 1); (1, 2), (1, 2), (1, 2), (1, 2))$	$(1, 1, (1, 1);)$
2	$(2, 1, (1, 1);)$	$(1, 1, (1, 1);)$
3	$(1, 2, (1, 1);)$	$(3, 0, (1, 1); (1, 3))$
3	$(3, 0, (1, 1); (2, 3), (2, 3))$	$(1, 1, (1, 1);)$
3	$(3, 0, (2, 2); (1, 3), (1, 3))$	$(3, 0, (2, 2); (1, 3), (1, 3))$
4	$(1, 2, (1, 1);)$	$(4, 0, (1, 1); (1, 2))$
5	$(5, 0, (1, 1); (3, 5))$	$(1, 1, (1, 1);)$
6	$(6, 0, (1, 1); (2, 3))$	$(1, 1, (1, 1);)$
6	$(2, 0, (1, 1); (1, 2), (1, 2), (1, 2), (1, 2))$	$(3, 0, (2, 2); (2, 3))$
6	$(2, 0, (1, 1); (1, 2), (1, 2), (1, 2), (1, 2))$	$(3, 0, (2, 2); (2, 3))$
6	$(2, 1, (1, 1);)$	$(3, 0, (2, 2); (2, 3))$
6	$(3, 0, (2, 2); (1, 3), (1, 3))$	$(2, 0, (1, 1); (1, 2), (1, 2))$
6	$(6, 0, (5, 5); (1, 3))$	$(3, 0, (1, 1); (1, 3))$
12	$(3, 0, (1, 1); (2, 3), (2, 3))$	$(4, 0, (3, 3); (1, 2))$
12	$(6, 0, (5, 5); (1, 3))$	$(4, 0, (1, 1); (1, 2))$
15	$(5, 0, (3, 3); (4, 5))$	$(3, 0, (2, 2); (2, 3))$
20	$(5, 0, (4, 4); (2, 5))$	$(4, 0, (1, 1); (1, 2))$

Table 30: $\mathcal{C} = \mathcal{C}^{(2)}$, $S_4 = S_2 \#_{\mathcal{C}} S_1$ and $(r, k) = (2, 0)$

Degree	D_1	D_2	D_3
6	$(3, 0, 2; (2, 3), (2, 3))$	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$	$(6, 0; (1, 6), (1, 2), (1, 3))$
6	$(1, 1, 1;)$	$(6, 0, 1; (1, 2), (1, 3))$	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2))$

Table 31: $\mathcal{C} = \mathcal{C}^{(2)} \sqcup \{c_1\}$, $S_4 = S_1 \#_{c_1} S_1 \#_{\mathcal{C}^{(2)}} S_1$ and $(r, k) = (1, 1)$.

Degree	D_1	D_2	D_3
3	$(3, 0, 2; (2, 3), (2, 3))$	$(3, 0, (2, 2, 2);)$	$(3, 0, (2, 2); (2, 3))$
12	$(4, 0, 3; (1, 2), (3, 4))$	$(3, 0, (1, 1, 1);)$	$(4, 0, (3, 3); (1, 2))$

Table 32: $\mathcal{C} = \{c_1\} \sqcup \mathcal{C}^{(2)}$, $S_4 = S_1 \#_{c_1} S_1 \#_{\mathcal{C}^{(2)}} S_1$ and $(r, k) = (3, 0)$.

$\mathcal{C}$ is a mixed multicurve.

Degree	D_1	D_2
4	$(4, 0; (1, 4), (1, 4))$	$(1, 0;)$

Table 33: $\mathcal{C} = \{c_1, \dots, c_4, d_1, \dots, d_4\}$, $S_4 = S_0 \#_{\{c_1, \dots, c_4\}} S_1(4)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (0, 2)$.

Degree	D_1	D_2	D_3
2	$(1, 0;)$	$(2, 0; (1, 2), (1, 2))$	$(1, 0;)$

Table 34: $\mathcal{C} = \{c_1, \dots, c_4, d_1, \dots, d_4\}$, $S_4 = S_1(2) \#_{\{c_1, c_2\}} S_0 \#_{\{c_3, c_4\}} S_1(2)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (0, 4)$.

Degree	D_1	D_2	D_3
2	$(1, 1;)$	$(2, 0; (1, 2), (1, 2))$	$(1, 0;)$

Table 35: $\mathcal{C} = \{c_1, \dots, c_4, d_1, d_2\}$, $S_4 = S_1(2) \#_{\{c_1, c_2\}} S_0 \#_{\{c_3, c_4\}} S_1(2)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (0, 3)$

Degree	D_1	D_2	D_3	D_4
2	$(1, 0;)$	$(1, 1, 1;)$	$(2, 0, 1; (1, 2))$	$(1, 1, 1;)$

Table 36: $\mathcal{C} = \{c_1, \dots, c_4, d_1, d_2\}$, $S_4 = S_1(2) \#_{\{c_1, c_2\}} (S_1 \#_{c_3} S_0 \#_{c_4} S_1)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (2, 2)$

Degree	D_1	D_2
3	$(3, 1;)$	$(1, 0;)$

Table 37: $\mathcal{C} = \{c_1, c_2, c_3, d_1, d_2, d_3\}$, $S_4 = S_1 \#_{\{c_1, c_2, c_3\}} S_1(3)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (0, 2)$

Degree	D_1	D_2	D_3
2	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$	$(1, 1, 1;)$	$(4, 0; (1, 4), (1, 4))$

Table 38: $\mathcal{C} = \{c_1, c_2, c_3, d_1, d_2\}$, $S_4 = S_1 \#_{c_1} S_1 \#_{\{c_2, c_3\}} S_1(2)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (1, 2)$

Degree	D_1	D_2
2	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2), (1, 2))$	$(4, 0; (1, 4), (1, 4))$
2	$(2, 1; (1, 2), (1, 2))$	$(4, 0; (1, 4), (1, 4))$

Table 39: $\mathcal{C} = \{c_1, c_2, d_1, d_2\}$, $S_4 = S_2 \#_{\{c_1, c_2\}} S_1(2)$ with $d_i \in \widehat{S}_1^i$ and $(r, k) = (0, 2)$

Degree	D_1	D_2
2	$(2, 0; (1, 2), (1, 2))$	$(4, 0; (1, 4), (1, 2), (1, 4))$
2	$(2, 0; (1, 2), (1, 2))$	$(4, 0; (3, 4), (1, 2), (3, 4))$

Table 40: $\mathcal{C} = \{c_1, c_2, d_1, d_2\}$, $S_4 = S_2 \#_{\{c_1, c_2\}} S_1(2)$ with $d_i \in \widehat{S}_2$ and $(r, k) = (0, 2)$

Degree	D_1	D_2
2	$(1, 1, 1;)$	$(2, 0, 1; (1, 2), (1, 2), (1, 2))$

Table 41: $\mathcal{C} = \{c_1, d_1, d_2\}$, $S_4 = S_1 \#_{c_1} S_3$ with $\{d_1, d_2\} \subset \widehat{S}_3$ and $(r, k) = (1, 1)$

Degree	D_1	D_2
2	$(2, 0; (1, 2), (1, 2))$	$(2, 0; (1, 2), (1, 2), (1, 2), (1, 2))$

Table 42: $\mathcal{C} = \mathcal{C}^{(2)} \sqcup \{d_1, d_2\}$, $S_4 = S_2 \#_{\mathcal{C}^{(2)}} S_1$ with $\{d_1, d_2\} \subset \widehat{S}_2$ and $(r, k) = (0, 2)$.

8. CONCLUDING REMARKS

8.1. Roots and the Torelli Group. Let $\Psi : \text{Mod}(S_g) \rightarrow \text{Sp}(2g, \mathbb{Z})$ be the symplectic representation of $\text{Mod}(S_g)$ arising out of its action on $H_1(S_g, \mathbb{Z})$, and let $\mathcal{I}(S_g)$ denote the kernel of Ψ , the Torelli group of S_g . If $\mathcal{C}$ contains nonseparating curves, then $\Psi(t_{\mathcal{C}})$ is a product of commuting elementary matrices, and thus a root of $t_{\mathcal{C}}$ induces a root of such a matrix in $\text{Sp}(2g, \mathbb{Z})$. However, if every curve in $\mathcal{C}$ is separating, then $\Psi(h)$ is a root of unity in $\text{Sp}(2g, \mathbb{Z})$. Using the theory we have developed for multicurves, we shall now give a succinct proof of the fact that such a root cannot lie in $\mathcal{I}(S_g)$.

Theorem 8.1. *Let h be the root of the Dehn twist $t_{\mathcal{C}}$ about a multicurve $\mathcal{C}$ in S_g . Then $h \notin \mathcal{I}(S_g)$.*

Proof. Since $\mathcal{I}(S_g)$ is normal in $\text{Mod}(S_g)$, it suffices to prove that the conjugacy class of h intersects $\mathcal{I}(S_g)$ non-trivially. If c is an essential simple closed curve in S_g , it follows from [3, Proposition 6.3, Example 6.5.2] that $t_c \in \mathcal{I}(S_g)$ if and only if c is a separating curve. Consequently, it suffices to assume that every curve in $\mathcal{C}$ is a separating curve.

In that case, let $\text{Fix}_h(\mathcal{C}) = \{\mathcal{E}_1, \dots, \mathcal{E}_m\}$, where $m = |\text{Fix}_h(\mathcal{C})|$, and write

$$S_g = \overline{\#}_{i=1}^{m(h)} (S_{g_i} \#_{\mathcal{E}_i} S_{g_{i+1}}),$$

as in Notation 5.14. By Theorem 5.17, the conjugacy class of h corresponds to an $(m+1)$ -compatible multituple $(\overline{t}_1, \overline{t}_2, \dots, \overline{t}_{m+1})$, where $\overline{t}_i = (\llbracket t_{i,1} \rrbracket, \llbracket t_{i,2,1} \rrbracket, \dots, \llbracket t_{i,2,s_i} \rrbracket)$ is a (s_i+1) -compatible tuple of genus g_i .

Choose $1 \leq i \leq m$ such that at least one component of $\overline{t}_i$ is non-trivial, so we write

$$S_{g_i} = \overline{\#}_{j=1}^{k_i} (S_{g_{i,1}} \#_{D_{i,j}} \mathbb{S}_{g_{i,2,j}}(m_{1,j})),$$

as in Notation 5.14. If $g_{i,1} \geq 1$, and $t_{i,1}$ is a non-trivial finite order element in $\text{Mod}(S_{g_{i,1}})$, then $t_{i,1} \notin \mathcal{I}(S_{g_{i,1}})$ by [3, Theorem 6.8]. Since $t_{i,1}$ is obtained from a restriction of h by coning, it follows that $h \notin \mathcal{I}(S_g)$. If $t_{i,1}$ is trivial or $g_{i,1} = 0$, then it follows $k_i \geq 1$ and that $t_{i,2,j}$ is non-trivial for some $1 \leq j \leq k_i$. Since each curve in $\mathcal{C}$ is separating, $g_{i,2,j} \geq 1$, and so the surface $\mathbb{S}_{g_{i,2,j}}(m_{1,j})$ contributes $2m_{1,j}g_{i,2,j}$ generators to the standard geometric symplectic basis of $H_1(S_g, \mathbb{Z})$. Now $t_{i,2,j}$ (and consequently h) cyclically permutes the components of $\mathbb{S}_{g_{i,2,j}}(m_{1,j})$, and thus $h \notin \mathcal{I}(S_g)$. $\square$

8.2. Roots of finite product of powers. The theory developed in this paper for classifying roots up to conjugacy for finite products of commuting Dehn twists can be naturally generalised to one that classifies roots of finite products of powers of commuting twists.

Currently, the compatibility condition requires that pairs of distinguished orbits (or fixed points) of permuting actions should have associated angles that add up to $2\pi/n \pmod{2\pi}$. When c is a single nonseparating curve, the roots of t_c^ℓ for $1 \leq \ell < n$, were classified in [10] by using a variant of this condition, which required that the angles associated with compatible fixed points add up to $2\pi\ell/n \pmod{2\pi}$. This notion of compatibility of fixed points can be generalized to orbits, and this will eventually lead to the classification of roots of homeomorphisms of the form $\prod_{i=1}^m t_{c_i}^{\ell_i}$, where $\{c_1, c_2, \dots, c_m\}$ is a multicurve. It is also apparent that such roots would not lie in $\mathcal{I}(S_g)$ for the same reasons as above.

9. ACKNOWLEDGEMENTS

The authors would like to thank Dan Margalit and Darryl McCullough for helpful suggestions and examples that made the theory more comprehensive.

REFERENCES

- [1] Robert J Daverman and Richard B Sher. *Handbook of geometric topology*. North-Holland, 2002.
- [2] Allan L. Edmonds. Surface symmetry. I. *Michigan Math. J.*, 29(2):171–183, 1982.
- [3] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [4] W. J. Harvey. Cyclic groups of automorphisms of a compact Riemann surface. *Quart. J. Math. Oxford Ser. (2)*, 17:86–97, 1966.
- [5] Steven P. Kerckhoff. The Nielsen realization problem. *Bull. Amer. Math. Soc. (N.S.)*, 2(3):452–454, 1980.
- [6] Dan Margalit and Saul Schleimer. Dehn twists have roots. *Geom. Topol.*, 13(3):1495–1497, 2009.
- [7] Darryl McCullough and Kashyap Rajeevsarathy. Roots of Dehn twists. *Geometriae Dedicata*, 151:397–409, 2011. 10.1007/s10711-010-9541-4.
- [8] Jakob Nielsen. Abbildungsklassen endlicher Ordnung. *Acta Math.*, 75:23–115, 1943.
- [9] Kashyap Rajeevsarathy. Roots of dehn twists about separating curves. *Journal of the Australian Mathematical Society*, 95:266–288, 10 2013.
- [10] Kashyap Rajeevsarathy. Fractional powers of Dehn twists about nonseparating curves. *Glasg. Math. J.*, 56(1):197–210, 2014.
- [11] William P. Thurston. *The Geometry and Topology of Three-Manifolds*. notes available at: <http://www.msri.org/communications/books/gt3m/PDF>.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH BHOPAL, BHOPAL BYPASS ROAD, BHAURI, BHOPAL 462 066, MADHYA PRADESH, INDIA

E-mail address: kashyap@iiserb.ac.in

URL: <https://home.iiserb.ac.in/~kashyap/>

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH BHOPAL, BHOPAL BYPASS ROAD, BHAURI, BHOPAL 462 066, MADHYA PRADESH, INDIA

E-mail address: prahlad@iiserb.ac.in

URL: <https://home.iiserb.ac.in/~prahlad/>